

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
25 March 2004 (25.03.2004)

PCT

(10) International Publication Number  
**WO 2004/024322 A2**

- (51) International Patent Classification<sup>7</sup>: **B01J 31/24**, C07C 67/38 (74) Agents: **FRITH, Richard, William** et al.; Appleyard Lees, 15 Clare Road, Halifax HX1 2HY (GB).
- (21) International Application Number: PCT/GB2003/003936 (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (22) International Filing Date: 10 September 2003 (10.09.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
- |                 |                                |    |
|-----------------|--------------------------------|----|
| 0221093.8       | 12 September 2002 (12.09.2002) | GB |
| 0228018.8       | 30 November 2002 (30.11.2002)  | GB |
| 0309812.6       | 30 April 2003 (30.04.2003)     | GB |
| 0316159.3       | 10 July 2003 (10.07.2003)      | GB |
| PCT/GB03/003419 | 6 August 2003 (06.08.2003)     | GB |
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): **LUCITE INTERNATIONAL UK LIMITED** [GB/GB]; Wilton Centre, Wilton, Redcar TS10 4RF (GB).
- (72) Inventor; and
- (75) Inventor/Applicant (*for US only*): **EASTHAM, Graham** [GB/GB]; Wilton Centre, Wilton, Redcar TS10 4RF (GB).
- Published:**  
— *without international search report and to be republished upon receipt of that report*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: A CATALYST SYSTEM

(57) Abstract: A catalyst suitable for carbonylating ethylenically unsaturated compounds comprising a Group VIIIB metal or compound thereof and a metallocene.



**WO 2004/024322 A2**

### A CATALYST SYSTEM


The present invention relates to compounds, in particular compounds for catalysing the carbonylation of ethylenically unsaturated compounds, methods of preparing such compounds and the use of such compounds for catalysing the carbonylation of ethylenically unsaturated compounds.

10 The carbonylation of ethylenically unsaturated compounds using carbon monoxide in the presence of a catalyst system and an alcohol or water to yield the corresponding ester or carboxylic acid, respectively, is well known. Suitable catalyst systems comprise a Group VIII metal (e.g. palladium) and a phosphine ligand (e.g. an alkyl phosphine or a bidentate phosphine ligand as disclosed in WO-A-9619434).

Although catalyst systems have been developed which exhibit reasonable stability during the carbonylation process and permit relatively high reaction rates to be achieved, there still exists a need for improved catalyst systems. Suitably, the present invention aims to provide an improved catalyst for carbonylating ethylenically unsaturated compounds.

According to a first aspect, the present invention provides a catalyst suitable for carbonylating an ethylenically unsaturated compound, which catalyst is obtainable by combining:

- (a) a Group VIIIB metal or a compound thereof; and,
- (b) a compound of formula I or salt thereof:



wherein:

5     $A_1$  and  $A_2$ , and  $A_3$ ,  $A_4$  and  $A_5$  (when present), each independently represent lower alkylene;


$K$  is selected from the group consisting of hydrogen, lower alkyl, aryl, Het, halo, cyano, nitro,  $-OR^{19}$ ,  $-OC(O)R^{20}$ ,  
 10     $-C(O)R^{21}$ ,  $-C(O)OR^{22}$ ,  $-N(R^{23})R^{24}$ ,  $-C(O)N(R^{25})R^{26}$ ,  $-C(S)(R^{27})R^{28}$ ,  
        $-SR^{29}$ ,  $-C(O)SR^{30}$ ,  $-CF_3$  or  $-A_3-Q^3(X^5)X^6$ ;

$D$  is selected from the group consisting of hydrogen, lower alkyl, aryl, Het, halo, cyano, nitro,  $-OR^{19}$ ,  $-OC(O)R^{20}$ ,  
 15     $-C(O)R^{21}$ ,  $-C(O)OR^{22}$ ,  $-N(R^{23})R^{24}$ ,  $-C(O)N(R^{25})R^{26}$ ,  $-C(S)(R^{27})R^{28}$ ,  
        $-SR^{29}$ ,  $-C(O)SR^{30}$ ,  $-CF_3$  or  $-A_4-Q^4(X^7)X^8$ ;

$E$  is selected from the group consisting of hydrogen, lower alkyl, aryl, Het, halo, cyano, nitro,  $-OR^{19}$ ,  $-OC(O)R^{20}$ ,  
 20     $-C(O)R^{21}$ ,  $-C(O)OR^{22}$ ,  $-N(R^{23})R^{24}$ ,  $-C(O)N(R^{25})R^{26}$ ,  $-C(S)(R^{27})R^{28}$ ,  
        $-SR^{29}$ ,  $-C(O)SR^{30}$ ,  $-CF_3$  or  $-A_5-Q^5(X^9)X^{10}$ ;


or both D and E together with the carbon atoms of the cyclopentadienyl ring to which they are attached form an optionally substituted phenyl ring:

5  $X^1$  represents  $CR^1(R^2)(R^3)$ , congressyl or adamantyl,  $X^2$  represents  $CR^4(R^5)(R^6)$ , congressyl or adamantyl, or  $X^1$  and  $X^2$  together with  $Q^2$  to which they are attached form an optionally substituted 2-phospha-  
 tricyclo[3.3.1.1{3,7}]decyl group or derivative thereof,  
 10 or  $X^1$  and  $X^2$  together with  $Q^2$  to which they are attached form a ring system of formula 1a



(1a)


$X^3$  represents  $CR^7(R^8)(R^9)$ , congressyl or adamantyl,  $X^4$  represents  $CR^{10}(R^{11})(R^{12})$ , congressyl or adamantyl, or  $X^3$  and  $X^4$  together with  $Q^1$  to which they are attached form an optionally substituted 2-phospha-  
 tricyclo[3.3.1.1{3,7}]decyl group or derivative thereof,  
 15 or  $X^3$  and  $X^4$  together with  $Q^1$  to which they are attached form a ring system of formula 1b



(1b)


$X^5$  represents  $CR^{13}(R^{14})(R^{15})$ , congressyl or adamantyl,  $X^6$  represents  $CR^{16}(R^{17})(R^{18})$ , congressyl or adamantyl, or  $X^5$

and  $X^6$  together with  $Q^3$  to which they are attached form an optionally substituted 2-phosphatricyclo[3.3.1.1{3,7}]decyl group or derivative thereof, or  $X^5$  and  $X^6$  together with  $Q^3$  to which they are attached  
 5 form a ring system of formula 1c




(1c)

$X^7$  represents  $CR^{31}(R^{32})(R^{33})$ , congressyl or adamantyl,  $X^8$  represents  $CR^{34}(R^{35})(R^{36})$ , congressyl or adamantyl, or  $X^7$  and  $X^8$  together with  $Q^4$  to which they are attached form an optionally substituted 2-phosphatricyclo[3.3.1.1{3,7}]decyl group or derivative thereof, or  $X^7$  and  $X^8$  together with  $Q^4$  to which they are attached  
 10 form a ring system of formula 1d



(1d)

$X^9$  represents  $CR^{37}(R^{38})(R^{39})$ , congressyl or adamantyl,  $X^{10}$  represents  $CR^{40}(R^{41})(R^{42})$ , congressyl or adamantyl, or  $X^9$  and  $X^{10}$  together with  $Q^5$  to which they are attached form an optionally substituted 2-phosphatricyclo[3.3.1.1{3,7}]decyl group or derivative thereof, or  $X^9$  and  $X^{10}$  together with  $Q^5$  to which they are attached  
 20 form a ring system of formula 1e



$Q^1$  and  $Q^2$ , and  $Q^3$ ,  $Q^4$  and  $Q^5$  (when present), each independently represent phosphorus, arsenic or antimony;

5

M represents a Group VIB or VIIIB metal or metal cation thereof;

$L_1$  represents an optionally substituted cyclopentadienyl, indenyl or aryl group;

10

$L_2$  represents one or more ligands each of which are independently selected from hydrogen, lower alkyl, alkylaryl, halo, CO,  $P(R^{43})(R^{44})R^{45}$  or  $N(R^{46})(R^{47})R^{48}$ ;

15

$R^1$  to  $R^{18}$  and  $R^{31}$  to  $R^{42}$ , when present, each independently represent hydrogen, lower alkyl, aryl, halo or Het;

$R^{19}$  to  $R^{30}$  and  $R^{43}$  to  $R^{48}$ , when present, each independently represent hydrogen, lower alkyl, aryl or Het;

20

$R^{49}$ ,  $R^{54}$  and  $R^{55}$ , when present, each independently represent hydrogen, lower alkyl or aryl;

$R^{50}$  to  $R^{53}$ , when present, each independently represent hydrogen, lower alkyl, aryl or Het;

25

$Y^1$ ,  $Y^2$ ,  $Y^3$ ,  $Y^4$  and  $Y^5$ , when present, each independently represent oxygen, sulfur or  $N-R^{55}$ ;

$n = 0$  or  $1$ ;

5

and  $m = 0$  to  $5$ ;

provided that when  $n = 1$  then  $m$  equals  $0$ , and when  $n$  equals  $0$  then  $m$  does not equal  $0$ .

10

Such compounds are referred to hereinafter as "the compounds of the invention".

Preferably in a compound of formula I when both K represents  $-A_3-Q^3(X^5)X^6$  and E represents  $-A_5-Q^5(X^9)X^{10}$ , then D represents  $-A_4-Q^4(X^7)X^8$ .

Suitably, the compounds of the invention may catalyse the carbonylation of unsaturated compounds, particularly ethylenically unsaturated compounds, in the presence of carbon monoxide and a coreactant. In particular, the compounds of the invention may be employed in hydroformylation reactions, hydrocarboxylation reactions, hydroesterification reactions and hydroamidation reactions. For example, the compounds of the invention may catalyse the conversion of an ethylenically unsaturated compound in the presence of carbon monoxide and a hydroxyl group containing compound to the corresponding carboxylic acid or ester, respectively, depending on the choice of hydroxyl group containing compound used (i.e. a carboxylic acid if the hydroxyl group containing compound is water and an ester if the hydroxyl group containing compound is an alcohol.

Conveniently, the compounds of the invention may exhibit a high stability under typical carbonylation reaction conditions such that they require little or no replenishment. Conveniently, the compounds of the invention may increase the rate of the carbonylation reaction of an ethylenically unsaturated compound compared to known catalysts. Conveniently, the compounds of the invention may promote high conversion rates of the ethylenically unsaturated compound, thereby yielding the desired product in high yield with little or no impurities. Consequently, employing the compounds of the invention may increase the commercial viability of a carbonylation process, such as the carbonylation of an ethylenically unsaturated compound. Suitably, the compounds of the invention typically exhibit a high catalytic turnover in the carbonylation of an unsaturated compound, particularly an ethylenically unsaturated compound, in the presence of carbon monoxide and a co-reactant as defined herein.

Unexpectedly, it has been found that if the carbon atoms of  $X^1$  and  $X^2$  bonded to  $Q^2$  and the carbon atoms of  $X^3$  and  $X^4$  bonded to  $Q^1$  in the compound of formula I do not include a hydrogen atom, then this typically produces a catalyst which exhibits increased catalytic turnover in the carbonylation of an unsaturated compound compared with a comparable compound where a carbon atom present in  $X^1$  and/or  $X^2$ , and  $X^3$  and/or  $X^4$  bonded to  $Q^2$  and  $Q^1$ , respectively, includes a hydrogen atom. Most preferably,  $Q^1$  is bonded to a tertiary carbon atom of both  $X^3$  and  $X^4$ , and  $Q^2$  is bonded to a tertiary carbon atom of both  $X^1$  and  $X^2$ . Similarly, if present, the carbon atoms of  $X^5$  and  $X^6$ ,



bonded to Q<sup>3</sup>, of X<sup>7</sup> and X<sup>8</sup> bonded to Q<sup>4</sup>, and of X<sup>9</sup> and X<sup>10</sup> bonded to Q<sup>5</sup> do not include a hydrogen atom. More preferably, Q<sup>3</sup>, Q<sup>4</sup> and Q<sup>5</sup>, if present, are bonded to a tertiary carbon atom of both X<sup>5</sup> and X<sup>6</sup>, X<sup>7</sup> and X<sup>8</sup>, and X<sup>9</sup> and X<sup>10</sup>, respectively.

Preferably, R<sup>1</sup> to R<sup>18</sup> and R<sup>31</sup> to R<sup>42</sup>, when present, each independently represent hydrogen, optionally substituted C<sub>1</sub> to C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkyl phenyl (wherein the phenyl group is optionally substituted as defined herein), trifluoromethyl or phenyl (wherein the phenyl group is optionally substituted as defined herein). Even more preferably, R<sup>1</sup> to R<sup>18</sup> and R<sup>31</sup> to R<sup>42</sup>, when present, each independently represent hydrogen, C<sub>1</sub> to C<sub>6</sub> alkyl, which is optionally substituted as defined herein, trifluoromethyl or optionally substituted phenyl. Even more preferably, R<sup>1</sup> to R<sup>18</sup> and R<sup>31</sup> to R<sup>42</sup>, when present each independently represent hydrogen, non-substituted C<sub>1</sub> to C<sub>6</sub> alkyl or phenyl which is optionally substituted with one or more substituents selected from non-substituted C<sub>1</sub> to C<sub>6</sub> alkyl or OR<sup>19</sup> where R<sup>19</sup> represents hydrogen or unsubstituted C<sub>1</sub> to C<sub>6</sub> alkyl. More preferably, R<sup>1</sup> to R<sup>18</sup> and R<sup>31</sup> to R<sup>42</sup>, when present, each independently represent hydrogen or non-substituted C<sub>1</sub> to C<sub>6</sub> alkyl such as methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, pentyl, hexyl and cyclohexyl, especially methyl. Most preferably, R<sup>1</sup> to R<sup>18</sup> and R<sup>31</sup> to R<sup>42</sup> when present, each independently represent non-substituted C<sub>1</sub> to C<sub>6</sub> alkyl such as methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, pentyl, hexyl and cyclohexyl, especially methyl.

Alternatively, or additionally, one or more of the groups R<sup>1</sup> to R<sup>3</sup>, R<sup>4</sup> to R<sup>6</sup>, R<sup>7</sup> to R<sup>9</sup>, R<sup>10</sup> to R<sup>12</sup>, R<sup>13</sup> to R<sup>15</sup>, R<sup>16</sup> to

$R^{18}$ ,  $R^{31}$  to  $R^{33}$ ,  $R^{34}$  to  $R^{36}$ ,  $R^{37}$  to  $R^{39}$  or  $R^{40}$  to  $R^{42}$  (when present) together with the carbon atom to which they are attached independently may form cyclic alkyl structures such as 1-norbornyl or 1-norbornadienyl.

5

Alternatively, or additionally, one or more of the groups  $R^1$  and  $R^2$ ,  $R^4$  and  $R^5$ ,  $R^7$  and  $R^8$ ,  $R^{10}$  and  $R^{11}$ ,  $R^{13}$  and  $R^{14}$ ,  $R^{16}$  and  $R^{17}$ ,  $R^{31}$  and  $R^{32}$ ,  $R^{34}$  and  $R^{35}$ ,  $R^{37}$  and  $R^{38}$  or  $R^{40}$  and  $R^{41}$  (when present) together with the carbon atom to which they are attached independently may form a cyclic alkyl structures, preferably a  $C_5$  to  $C_7$  cyclic alkyl structure such as cyclohexyl and cyclopentyl, and  $R^3$ ,  $R^6$ ,  $R^9$ ,  $R^{12}$ ,  $R^{15}$ ,  $R^{18}$ ,  $R^{33}$ ,  $R^{36}$ ,  $R^{39}$  and  $R^{42}$  (when present) each independently represent hydrogen, lower alkyl, trifluoromethyl or aryl as defined above, particularly non-substituted  $C_1$  to  $C_6$  alkyl and hydrogen, especially non-substituted  $C_1$  to  $C_6$  alkyl.

In an especially preferred embodiment, each of  $R^1$  to  $R^{18}$  and  $R^{31}$  to  $R^{42}$ , when present, do not represent hydrogen. Suitably, such an arrangement means  $Q^1$ ,  $Q^2$ ,  $Q^3$ ,  $Q^4$  and  $Q^5$  are bonded to a carbon atom of  $X^1$  to  $X^{10}$ , respectively, which bears no hydrogen atoms.

Preferably,  $R^1$ ,  $R^4$ ,  $R^7$ ,  $R^{10}$ ,  $R^{13}$ ,  $R^{16}$ ,  $R^{31}$ ,  $R^{34}$ ,  $R^{37}$  and  $R^{40}$  (when present), each represent the same substituent as defined herein;  $R^2$ ,  $R^5$ ,  $R^8$ ,  $R^{11}$ ,  $R^{14}$ ,  $R^{17}$ ,  $R^{32}$ ,  $R^{35}$ ,  $R^{38}$  and  $R^{41}$  (when present), each represent the same substituent as defined herein; and  $R^3$ ,  $R^6$ ,  $R^9$ ,  $R^{12}$ ,  $R^{15}$ ,  $R^{18}$ ,  $R^{33}$ ,  $R^{36}$ ,  $R^{39}$  and  $R^{42}$  (when present), each represent the same substituent as defined herein. More preferably  $R^1$ ,  $R^4$ ,  $R^7$ ,  $R^{10}$ ,  $R^{13}$ ,  $R^{16}$ ,  $R^{31}$ ,  $R^{34}$ ,  $R^{37}$  and  $R^{40}$  (when present) each represent the same  $C_1$ - $C_6$  alkyl, particularly non-

substituted C<sub>1</sub>-C<sub>6</sub> alkyl, such as methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, pentyl, hexyl or cyclohexyl, or trifluoromethyl; R<sup>2</sup>, R<sup>5</sup>, R<sup>8</sup>, R<sup>11</sup>, R<sup>14</sup>, R<sup>17</sup>, R<sup>32</sup>, R<sup>35</sup>, R<sup>38</sup> and R<sup>41</sup> (when present), each independently  
5 represent the same C<sub>1</sub>-C<sub>6</sub> alkyl as defined above, or trifluoromethyl; and R<sup>3</sup>, R<sup>6</sup>, R<sup>9</sup>, R<sup>12</sup>, R<sup>15</sup>, R<sup>18</sup>, R<sup>33</sup>, R<sup>36</sup>, R<sup>39</sup> and R<sup>42</sup> (when present), each independently represent the same C<sub>1</sub>-C<sub>6</sub> alkyl as defined above, or trifluoromethyl. For example: R<sup>1</sup>, R<sup>4</sup>, R<sup>7</sup>, R<sup>10</sup>, R<sup>13</sup> and R<sup>16</sup> (when present) each  
10 represent methyl; R<sup>2</sup>, R<sup>5</sup>, R<sup>8</sup>, R<sup>11</sup>, R<sup>14</sup> and R<sup>17</sup> each represent ethyl (when present); and, R<sup>3</sup>, R<sup>6</sup>, R<sup>9</sup>, R<sup>12</sup>, R<sup>15</sup> and R<sup>18</sup> (when present) each represent n-butyl or n-pentyl.

In an especially preferred embodiment of the present  
15 invention each R<sup>1</sup> to R<sup>18</sup> and R<sup>31</sup> to R<sup>42</sup> group (when present) represents the same substituent as defined herein. Preferably, each R<sup>1</sup> to R<sup>18</sup> and R<sup>31</sup> to R<sup>42</sup> group represents the same C<sub>1</sub> to C<sub>6</sub> alkyl group, particularly non-substituted C<sub>1</sub>-C<sub>6</sub> alkyl, such as methyl, ethyl, n-propyl,  
20 iso-propyl, n-butyl, iso-butyl, tert-butyl, pentyl, hexyl and cyclohexyl, or trifluoromethyl. Most preferably, each R<sup>1</sup> to R<sup>18</sup> and R<sup>31</sup> to R<sup>42</sup> group represents non-substituted C<sub>1</sub>-C<sub>6</sub> alkyl, particularly methyl.

25 The term adamantyl when used herein means an adamantyl group which may be bonded to Q<sup>1</sup>, Q<sup>2</sup>, Q<sup>3</sup>, Q<sup>4</sup> and Q<sup>5</sup>, respectively, in position 1 or 2. Tricyclo[3.3.1.1.{3,7}]decyl is the systematic name for an adamantyl group, suitably Q<sup>1</sup>, Q<sup>2</sup>, Q<sup>3</sup>, Q<sup>4</sup> and Q<sup>5</sup>,  
30 respectively, may be bonded to the 1 position or 2 position of one or two tricyclo[3.3.1.1.{3,7}]decyl groups. Preferably, Q<sup>1</sup> and Q<sup>2</sup>, and Q<sup>3</sup>, Q<sup>4</sup> and Q<sup>5</sup>, when present, is bonded to a tertiary carbon of one or more

adamantyl groups. Suitably, when the adamantyl group represents unsubstituted adamantyl,  $Q^1$  and  $Q^2$ , and  $Q^3$ ,  $Q^4$  and  $Q^5$  when present are preferably bonded to the 1 position of one or more tricyclo[3.3.1.1{3,7}]decyl groups  
5 i.e. the carbon atom of the adamantyl group bears no hydrogen atom.

The adamantyl group may optionally comprise, besides hydrogen atoms, one or more substituents selected from  
10 lower alkyl,  $-OR^{19}$ ,  $-OC(O)R^{20}$ , halo, nitro,  $-C(O)R^{21}$ ,  $-C(O)OR^{22}$ , cyano, aryl,  $-N(R^{23})R^{24}$ ,  $-C(O)N(R^{25})R^{26}$ ,  $-C(S)(R^{27})R^{28}$ ,  $-CF_3$ ,  $-P(R^{56})R^{57}$ ,  $-PO(R^{58})(R^{59})$ ,  $-PO_3H_2$ ,  $-PO(OR^{60})(OR^{61})$ , or  $-SO_3R^{62}$ , wherein  $R^{19}$ ,  $R^{20}$ ,  $R^{21}$ ,  $R^{22}$ ,  $R^{23}$ ,  $R^{24}$ ,  $R^{25}$ ,  $R^{26}$ ,  $R^{27}$ ,  $R^{28}$ , lower alkyl, cyano and aryl are as  
15 defined herein and  $R^{56}$  to  $R^{62}$  each independently represent hydrogen, lower alkyl, aryl or Het.

Suitably, when the adamantyl group is substituted with one or more substituents as defined above, highly preferred  
20 substituents include unsubstituted  $C_1$  to  $C_8$  alkyl,  $-OR^{19}$ ,  $-OC(O)R^{20}$ , phenyl,  $-C(O)OR^{22}$ , fluoro,  $-SO_3H$ ,  $-N(R^{23})R^{24}$ ,  $-P(R^{56})R^{57}$ ,  $-C(O)N(R^{25})R^{26}$  and  $-PO(R^{58})(R^{59})$ ,  $-CF_3$ , wherein  $R^{19}$  represents hydrogen, unsubstituted  $C_1$ - $C_8$  alkyl or phenyl,  $R^{20}$ ,  $R^{22}$ ,  $R^{23}$ ,  $R^{24}$ ,  $R^{25}$ ,  $R^{26}$  each independently represent  
25 hydrogen or unsubstituted  $C_1$ - $C_8$  alkyl,  $R^{56}$  to  $R^{59}$ ,  $R^{56}$  each independently represent unsubstituted  $C_1$ - $C_8$  alkyl or phenyl.

Suitably, the adamantyl group may comprise, besides  
30 hydrogen atoms, up to 10 substituents as defined above, preferably up to 5 substituents as defined above, more preferably up to 3 substituents as defined above. Suitably, when the adamantyl group comprises, besides

hydrogen atoms, one or more substituents as defined herein, preferably each substituent is identical. Preferred substituents are unsubstituted C<sub>1</sub>-C<sub>8</sub> alkyl and trifluoromethyl, particularly unsubstituted C<sub>1</sub>-C<sub>8</sub> alkyl  
5 such as methyl. A highly preferred adamantyl group comprises hydrogen atoms only i.e. the adamantyl group is not substituted.

Preferably, when more than one adamantyl group is present  
10 in a compound of formula I, each adamantyl group is identical.

By the term 2-phospha-tricyclo[3.3.1.1.{3,7}]decyl group we mean a 2-phospha-adamantyl group formed by the  
15 combination of X<sup>1</sup> and X<sup>2</sup> together with Q<sup>2</sup> to which they are attached, a 2-phospha-adamantyl group formed by the combination of X<sup>3</sup> and X<sup>4</sup> together with Q<sup>1</sup> to which they are attached, a 2-phospha-adamantyl group formed by the combination of X<sup>5</sup> and X<sup>6</sup> together with Q<sup>3</sup> to which they are  
20 attached, a 2-phospha-adamantyl group formed by the combination of X<sup>7</sup> and X<sup>8</sup> together with Q<sup>4</sup> to which they are attached and a 2-phospha-adamantyl group formed by the combination of X<sup>9</sup> and X<sup>10</sup> together with Q<sup>5</sup> to which they are attached, wherein Q<sup>1</sup>, Q<sup>2</sup>, Q<sup>3</sup>, Q<sup>4</sup> and Q<sup>5</sup> is in the 2-  
25 position of the adamantyl group of which it forms an integral part and each of Q<sup>1</sup>, Q<sup>2</sup>, Q<sup>3</sup>, Q<sup>4</sup> and Q<sup>5</sup> represents phosphorus.

The 2-phospha-tricyclo[3.3.1.1.{3,7}]decyl group (referred  
30 to as 2-phospha-adamantyl group herein) may optionally comprise, beside hydrogen atoms, one or more substituents. Suitable substituents include those substituents as defined herein in respect of the adamantyl group. Highly

preferred substituents include lower alkyl, particularly unsubstituted C<sub>1</sub>-C<sub>8</sub> alkyl, especially methyl, trifluoromethyl, -OR<sup>19</sup> wherein R<sup>19</sup> is as defined herein particularly unsubstituted C<sub>1</sub>-C<sub>8</sub> alkyl or aryl, and 4-  
5 dodecylphenyl. When the 2-phospha-adamantyl group includes more than one substituent, preferably each substituent is identical.

Preferably, the 2-phospha-adamantyl group is substituted  
10 on one or more of the 1, 3, 5 or 7 positions with a substituent as defined herein. More preferably, the 2-phospha-adamantyl group is substituted on each of the 1, 3 and 5 positions. Suitably, such an arrangement means the phosphorous atom of the 2-phospha-adamantyl group is  
15 bonded to carbon atoms in the adamantyl skeleton having no hydrogen atoms. Most preferably, the 2-phospha-adamantyl group is substituted on each of the 1, 3, 5 and 7 positions. When the 2-phospha-adamantyl group includes more than 1 substituent preferably each substituent is  
20 identical. Especially preferred substituents are unsubstituted C<sub>1</sub>-C<sub>8</sub> alkyl and trifluoromethyl, particularly unsubstituted C<sub>1</sub>-C<sub>8</sub> alkyl such as methyl.

Preferably, the 2-phospha-adamantyl group includes  
25 additional heteroatoms, other than the 2-phosphorous atom, in the 2-phospha-adamantyl skeleton. Suitable additional heteroatoms include oxygen and sulphur atoms, especially oxygen atoms. More preferably, the 2-phospha-adamantyl group includes one or more additional heteroatoms in the  
30 6, 9 and 10 positions. Even more preferably, the 2-phospha-adamantyl group includes an additional heteroatom in each of the 6, 9 and 10 positions. Most preferably, when the 2-phospha-adamantyl group includes two or more

additional heteroatoms in the 2-phospha-adamantyl skeleton, each of the additional heteroatoms are identical. An especially preferred 2-phospha-adamantyl group, which may optionally be substituted with one or more substituents as defined herein, includes an oxygen atom in each of the 6, 9 and 10 positions of the 2-phospha-adamantyl skeleton.

Highly preferred 2-phospha-adamantyl groups as defined herein include 2-phospha-1,3,5,7-tetramethyl-6,9,10-trioxadamantyl group, 2-phospha-1,3,5-trimethyl-6,9,10-trioxadamantyl group, 2-phospha-1,3,5,7-tetra(trifluoromethyl)-6,9,10-trioxadamantyl group, and 2-phospha-1,3,5-tri(trifluoromethyl)-6,9,10-trioxadamantyl group. Most preferably, the 2-phospha-adamantyl is selected from 2-phospha-1,3,5,7-tetramethyl-6,9,10-trioxadamantyl group or 2-phospha-1,3,5-trimethyl-6,9,10-trioxadamantyl group.

Preferably, when more than one 2-phospha-adamantyl group is present in a compound of formula I, each 2-phospha-adamantyl group is identical.

The term congressyl when used herein means a congressyl group (also known as diamantyl group) which may be bonded to  $Q^1$ ,  $Q^2$ ,  $Q^3$ ,  $Q^4$  and  $Q^5$  respectively. Preferably,  $Q^1$  and  $Q^2$ , and  $Q^3$ ,  $Q^4$  and  $Q^5$ , when present, is bonded to one of the tertiary carbon atoms of the congressyl groups. Suitably, when the congressyl group is unsubstituted,  $Q^1$  and  $Q^2$ , and  $Q^3$ ,  $Q^4$  and  $Q^5$  when present, are preferably bonded to the 1-position of one or more congressyl groups.

The congressyl group may optionally comprise, beside hydrogen atoms, one or more substituents. Suitable substituents include those substituents as defined herein in respect of the adamantyl group. Highly preferred substituents include unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl groups, particularly methyl, and trifluoromethyl. Most preferably, the congressyl group is unsubstituted and comprises hydrogen atoms only.

10 Preferably, when more than one congressyl group is present in a compound of formula I, each congressyl group is identical.

Preferably, where one or more ring systems of formula Ia, Ib, Ic, Id or Ie are present in a compound of formula I, R<sup>50</sup> to R<sup>53</sup> each independently represent lower alkyl, aryl or Het, which groups are optionally substituted and/or terminated as defined herein. Such an arrangement means Q<sup>2</sup>, Q<sup>1</sup>, Q<sup>3</sup>, Q<sup>4</sup> and Q<sup>5</sup> of the ring system of formula Ia to Ie, respectively, is not bonded to a carbon atom bearing a hydrogen atom. Even more preferably, R<sup>50</sup> to R<sup>53</sup> each independently represent optionally substituted C<sub>1</sub>-C<sub>6</sub> alkyl, preferably non-substituted C<sub>1</sub>-C<sub>6</sub> alkyl, phenyl optionally substituted with non-substituted C<sub>1</sub>-C<sub>6</sub> alkyl or OR<sup>19</sup> where R<sup>19</sup> represents non-substituted C<sub>1</sub>-C<sub>6</sub> alkyl, or trifluoromethyl. Even more preferably R<sup>50</sup> to R<sup>53</sup> each represent the same group as defined herein, particularly non-substituted C<sub>1</sub>-C<sub>6</sub> alkyl, especially methyl.

30 Preferably, where one or more ring system of formula Ia to Ie are present in a compound of formula I, R<sup>49</sup> and R<sup>54</sup> each independently represent optionally substituted C<sub>1</sub>-C<sub>6</sub> alkyl, preferably non-substituted C<sub>1</sub>-C<sub>6</sub> alkyl, phenyl



optionally substituted with non-substituted C<sub>1</sub>-C<sub>6</sub> alkyl or OR<sup>19</sup> where R<sup>19</sup> represents non-substituted C<sub>1</sub>-C<sub>6</sub> alkyl, trifluoromethyl or hydrogen. More preferably, R<sup>49</sup> and R<sup>54</sup> represent the same group as defined herein, especially  
5 hydrogen.

Preferably, where one or more ring systems of formula Ia to Ie are present in a compound of formula I, Y<sup>1</sup> to Y<sup>5</sup> are identical. Most preferably, each of Y<sup>1</sup> to Y<sup>5</sup> represents  
10 oxygen. Preferably, where more than one ring system of formula Ia to Ie is present in a compound of formula I, each such ring system is identical.

Preferred embodiments of the present invention include  
15 those wherein:

X<sup>1</sup> represents CR<sup>1</sup>(R<sup>2</sup>)(R<sup>3</sup>), X<sup>2</sup> represents CR<sup>4</sup>(R<sup>5</sup>)(R<sup>6</sup>), X<sup>3</sup> represents CR<sup>7</sup>(R<sup>8</sup>)(R<sup>9</sup>) and X<sup>4</sup> represents CR<sup>10</sup>(R<sup>11</sup>)(R<sup>12</sup>);

20 X<sup>1</sup> represents CR<sup>1</sup>(R<sup>2</sup>)(R<sup>3</sup>), X<sup>2</sup> represents adamantyl, X<sup>3</sup> represents CR<sup>7</sup>(R<sup>8</sup>)(R<sup>9</sup>) and X<sup>4</sup> represents adamantyl;

X<sup>1</sup> represents CR<sup>1</sup>(R<sup>2</sup>)(R<sup>3</sup>), X<sup>2</sup> represents congressyl, X<sup>3</sup> represents CR<sup>7</sup>(R<sup>8</sup>)(R<sup>9</sup>) and X<sup>4</sup> represents congressyl;

25

X<sup>1</sup> represents CR<sup>1</sup>(R<sup>2</sup>)(R<sup>3</sup>), X<sup>2</sup> represents CR<sup>4</sup>(R<sup>5</sup>)(R<sup>6</sup>), and X<sup>3</sup> and X<sup>4</sup> together with Q<sup>1</sup> to which they are attached form a ring system of formula Ib or a 2-phospha-adamantyl group;

30 X<sup>1</sup> represents CR<sup>1</sup>(R<sup>2</sup>)(R<sup>3</sup>), X<sup>2</sup> represents adamantyl, X<sup>3</sup> and X<sup>4</sup> together with Q<sup>1</sup> to which they are attached form a ring system of formula Ib or a 2-phospha-adamantyl group;

$X^1$  represents  $CR^1(R^2)(R^3)$ ,  $X^2$  represents congressyl,  $X^3$  and  $X^4$  together with  $Q^1$  to which they are attached form a ring system of formula Ib or a 2-phospha-adamantyl group;

5.  $X^1$  to  $X^4$  each independently represent adamantyl;

$X^1$  to  $X^4$  each independently represent congressyl;

$X^1$  and  $X^2$  each independently represent adamantyl and  $X^3$  and  
10  $X^4$  each independently represent congressyl;

$X^1$  and  $X^3$  independently represent adamantyl and  $X^2$  and  $X^4$   
independently represent congressyl;

15  $X^1$  and  $X^2$  independently represent adamantyl,  $X^3$  represents  $CR^7(R^8)(R^9)$  and  $X^4$  represents  $CR^{10}(R^{11})(R^{12})$ ;

$X^1$  and  $X^2$  independently represent congressyl,  $X^3$  represents  $CR^7(R^8)(R^9)$  and  $X^4$  represents  $CR^{10}(R^{11})(R^{12})$ ;

20

$X^1$  and  $X^2$  independently represent adamantyl, and  $X^3$  and  $X^4$   
together with  $Q^1$  to which they are attached form a ring  
system of formula Ib or a 2-phospha-adamantyl group;

25  $X^1$  and  $X^2$  independently represent congressyl, and  $X^3$  and  $X^4$   
together with  $Q^1$  to which they are attached form a ring  
system of formula Ib or a 2-phospha-adamantyl group;

$X^1$  and  $X^2$  together with  $Q^2$  to which they are attached form  
30 a ring system of formula Ia, and  $X^3$  and  $X^4$  together with  $Q^1$   
to which they are attached form a ring system of formula  
Ib;

$X^1$  and  $X^2$  together with  $Q^2$  to which they are attached form a 2-phospha-adamantyl group, and  $X^3$  and  $X^4$  together with  $Q^1$  to which they are attached form a 2-phospha-adamantyl group;

5

Highly preferred embodiments of the present invention include those wherein:

$X^1$  represents  $CR^1(R^2)(R^3)$ ,  $X^2$  represents  $CR^4(R^5)(R^6)$ ,  $X^3$   
10 represents  $CR^7(R^8)(R^9)$  and  $X^4$  represents  $CR^{10}(R^{11})(R^{12})$ ;

$X^1$  represents  $CR^1(R^2)(R^3)$ ,  $X^2$  represents adamantyl,  $X^3$   
represents  $CR^7(R^8)(R^9)$  and  $X^4$  represents adamantyl;

15  $X^1$  represents  $CR^1(R^2)(R^3)$ ,  $X^2$  represents congressyl,  $X^3$   
represents  $CR^7(R^8)(R^9)$  and  $X^4$  represents congressyl;

$X^1$  to  $X^4$  each independently represent adamantyl;

20  $X^1$  to  $X^4$  each independently represent congressyl;

$X^1$  and  $X^2$  together with  $Q^2$  to which they are attached form a ring system of formula Ia, and  $X^3$  and  $X^4$  together with  $Q^1$  to which they are attached form a ring system of formula  
25 Ib;

$X^1$  and  $X^2$  together with  $Q^2$  to which they are attached form a 2-phospha-adamantyl group, and  $X^3$  and  $X^4$  together with  $Q^1$  to which they are attached form a 2-phospha-adamantyl  
30 group;

Preferably in a compound of formula I,  $X^1$  is identical to  $X^3$  and  $X^2$  is identical to  $X^4$ . More preferably,  $X^1$  is

identical to  $X^3$  and  $X^5$ ,  $X^7$  and  $X^9$  when present, and  $X^2$  is identical to  $X^4$  and  $X^6$ ,  $X^8$  and  $X^{10}$  when present. Even more preferably,  $X^1$  to  $X^4$  are identical. Most preferably,  $X^1$  to  $X^4$  are identical to each of  $X^6$  to  $X^{10}$  when present.

5

Preferably, in the compound of formula I,  $X^1$  and  $X^2$  represent identical substituents,  $X^3$  and  $X^4$  represent identical substituents,  $X^5$  and  $X^6$  (when present) represent identical substituents,  $X^7$  and  $X^8$  (when present) represent identical substituents, and  $X^9$  and  $X^{10}$  (when present) represent identical substituents.

Preferably, in a compound of formula I, K represents  $-A_3-Q^3(X^5)X^6$ , hydrogen, lower alkyl,  $-CF_3$ , phenyl or lower alkyl phenyl. More preferably, K represents  $-A_3-Q^3(X^5)X^6$ , hydrogen, unsubstituted  $C_1-C_6$  alkyl, unsubstituted phenyl, trifluoromethyl or  $C_1-C_6$  alkyl phenyl.

In a particular preferred embodiment K in a compound of formula I represents hydrogen.

In an alternative embodiment where K does not represent hydrogen, K represents  $-A_3-Q^3(X^5)X^6$ . Preferably,  $X^5$  is identical to  $X^3$  or  $X^1$ , and  $X^6$  is identical to  $X^2$  or  $X^4$ . More preferably,  $X^5$  is identical to both  $X^3$  and  $X^1$ , and  $X^6$  is identical to both  $X^2$  and  $X^4$ . Even more preferably,  $-A_3-Q^3(X^5)X^6$  is identical to either  $-A_1-Q^2(X^1)X^2$  or  $-A_2-Q^1(X^3)X^4$ . Most preferably,  $-A_3-Q^3(X^5)X^6$  is identical to both  $-A_1-Q^2(X^1)X^2$  and  $-A_2-Q^1(X^3)X^4$ .

30

Most preferably, K represents hydrogen in a compound of formula I.

Preferably, in the compound of formula I, D represents -  
A<sub>4</sub>-Q<sup>4</sup>(X<sup>7</sup>)X<sup>8</sup>, hydrogen, lower alkyl, CF<sub>3</sub>, phenyl or lower  
alkylphenyl, and E represents -A<sub>5</sub>-Q<sup>5</sup>(X<sup>9</sup>)X<sup>10</sup>, hydrogen,  
lower alkyl, CF<sub>3</sub>, phenyl or lower alkylphenyl, or D and E  
5 together with the carbons of the cyclopentadienyl ring to  
which they are attached form an optionally substituted  
phenyl ring. More preferably, D represents -A<sub>4</sub>-Q<sup>4</sup>(X<sup>7</sup>)X<sup>8</sup>,  
hydrogen, phenyl, C<sub>1</sub>-C<sub>6</sub> alkylphenyl, unsubstituted C<sub>1</sub>-C<sub>6</sub>  
alkyl, such as methyl, ethyl, propyl, butyl, pentyl and  
10 hexyl, or CF<sub>3</sub>; E represents -A<sub>5</sub>-Q<sup>5</sup>(X<sup>9</sup>)X<sup>10</sup>, hydrogen, phenyl,  
C<sub>1</sub>-C<sub>6</sub> alkylphenyl, unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl such as  
methyl, ethyl, propyl, butyl, pentyl and hexyl, or -CF<sub>3</sub>;  
or both D and E together with the carbon atoms of the  
cyclopentadienyl ring to which they are attached form a  
15 phenyl ring which is optionally substituted with one or  
more groups selected from phenyl, C<sub>1</sub>-C<sub>6</sub> alkylphenyl,  
unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl or -CF<sub>3</sub>.

Suitably, when D and E together with the carbon atoms of  
20 the cyclopentadienyl ring to which they are attached form  
an optionally substituted phenyl ring, the metal M or  
cation thereof is attached to an indenyl ring system.

In a particular preferred embodiment, D in a compound of  
25 formula I, represents hydrogen.

In an alternative embodiment where D does not represent  
hydrogen, D represents -A<sub>4</sub>-Q<sup>4</sup>(X<sup>7</sup>)X<sup>8</sup>. Preferably X<sup>8</sup> is  
identical to X<sup>4</sup> or X<sup>2</sup>, and X<sup>7</sup> is identical to X<sup>1</sup> or X<sup>3</sup>.  
30 More preferably, X<sup>8</sup> is identical to both X<sup>4</sup> and X<sup>2</sup>, and X<sup>7</sup>  
is identical to X<sup>1</sup> and X<sup>3</sup>. Even more preferably, -A<sub>4</sub>-  
Q<sup>4</sup>(X<sup>7</sup>)X<sup>8</sup> is identical to either -A<sub>1</sub>-Q<sup>2</sup>(X<sup>1</sup>)X<sup>2</sup> or -A<sub>2</sub>-Q<sup>1</sup>(X<sup>3</sup>)X<sup>4</sup>.

Most preferably,  $-A_4-Q^4(X^7)X^8$  is identical to both  $-A_2-Q^1(X^3)X^4$ , and  $-A_3-Q^3(X^5)X^6$  if present.

In a particular preferred embodiment, E in a compound of  
5 formula I represents hydrogen.

In an alternative embodiment where E does not represent hydrogen, E represents  $-A_5-Q^5(X^9)X^{10}$ . Preferably  $X^{10}$  is identical to  $X^4$  or  $X^2$ , and  $X^9$  is identical to  $X^1$  or  $X^3$ .  
10 More preferably,  $X^{10}$  is identical to both  $X^4$  and  $X^2$ , and  $X^9$  is identical to  $X^1$  and  $X^3$ . Even more preferably,  $-A_5-Q^5(X^9)X^{10}$  is identical to either  $-A_1-Q^2(X^1)X^2$  or  $-A_2-Q^1(X^3)X^4$ . Most preferably,  $-A_5-Q^5(X^9)X^{10}$  is identical to both  $-A_1-Q^2(X^1)X^2$  and  $-A_2-Q^1(X^3)X^4$ , and  $-A_3-Q^3(X^5)X^6$  and  $-A_4-Q^4(X^7)X^8$  if  
15 present.

Preferably, in the compound of formula I, when D and E together with the carbon atoms of the cyclopentadienyl ring to which they are attached do not form an optionally  
20 substituted phenyl ring, each of K, D and E represent an identical substituent.

In an alternative preferred embodiment, D and E together with the carbons of the cyclopentadienyl ring to which they are attached form an unsubstituted phenyl ring.  
25

Highly preferred embodiments of compounds of formula I include those wherein:

30 K, D and E are identical substituents as defined herein, particularly where K, D and E represent hydrogen;

K represents hydrogen, and D and E together with the carbons of the cyclopentadienyl ring to which they are attached form an unsubstituted phenyl ring;

- 5 K represents  $-A_3-Q^3(X^5)X^6$  as defined herein and both D and E represent H;

K represents  $-A_3-Q^3(X^5)X^6$  as defined herein and D and E together with the carbon atoms of the cyclopentadienyl  
10 ring to which they are attached form an unsubstituted phenyl ring;

K represents  $-A_3-Q^3(X^5)X^6$ , D represents  $-A_4-Q^4(X^7)X^8$  and E represents  $-A_5-Q^5(X^9)X^{10}$ .

15

Especially preferred compounds of formula I include those where both D and E represent hydrogen or D and E together with the carbon atoms of the cyclopentadienyl ring to which they are attached form an unsubstituted phenyl ring,  
20 particularly those compounds where both D and E represent hydrogen.

Preferably, in the compound of formula I,  $A_1$  and  $A_2$ , and  $A_3$ ,  $A_4$  and  $A_5$  (when present), each independently represent  
25  $C_1$  to  $C_6$  alkylene which is optionally substituted as defined herein, for example with lower alkyl groups. Suitably,  $A_1$  and  $A_2$ , and  $A_3$ ,  $A_4$  and  $A_5$  (when present) may include a chiral carbon atom. Preferably, the lower alkylene groups which  $A_1$  to  $A_5$  may represent are non-  
30 substituted. A particular preferred lower alkylene, which  $A_1$  to  $A_5$  may independently represent, is  $-CH_2-$  or  $-C_2H_4-$ . Most preferably, each of  $A_1$  and  $A_2$ , and  $A_3$ ,  $A_4$  and  $A_5$  (when

present), represent the same lower alkylene as defined herein, particularly  $-\text{CH}_2-$ .

In the compound of formula I, preferably each  $\text{Q}^1$  and  $\text{Q}^2$ ,  
5 and  $\text{Q}^3$ ,  $\text{Q}^4$  and  $\text{Q}^5$  (when present) are the same. Most preferably, each  $\text{Q}^1$  and  $\text{Q}^2$ , and  $\text{Q}^3$ ,  $\text{Q}^4$  and  $\text{Q}^5$  (when present), represents phosphorus.

It will be appreciated by those skilled in the art that  
10 the compounds of formula I (referred to as (b) above) may function as ligands that coordinate with the Group VIIIB metal or compound thereof (referred to as (a) above) to form the compounds of the invention. Typically, the Group VIIIB metal or compound thereof (a) coordinates to the one  
15 or more phosphorus, arsenic and/or antimony atoms of the compound of formula I. It will be appreciated that the compounds of formula I may be referred to broadly as "metallocenes".

20 Suitably, when  $n = 1$  and  $\text{L}_1$  represents an optionally substituted cyclopentadienyl or indenyl group, the compounds of formula I may contain either two cyclopentadienyl rings, two indenyl rings or one indenyl and one cyclopentadienyl ring (each of which ring systems  
25 may optionally be substituted as described herein). Such compounds may be referred to as "sandwich compounds" as the metal M or metal cation thereof is sandwiched by the two ring systems. The respective cyclopentadienyl and/or indenyl ring systems may be substantially coplanar with  
30 respect to each other or they may be tilted with respect to each other (commonly referred to as bent metallocenes).



Alternatively, when  $n = 1$  and  $L_1$  represents aryl, the compounds of the invention may contain either one cyclopentadienyl or one indenyl ring (each of which ring systems may optionally be substituted as described herein) and one aryl ring which is optionally substituted as defined herein. Suitably, when  $n = 1$  and  $L_1$  represents aryl then the metal  $M$  of the compounds of formula I as defined herein is typically in the form of the metal cation.

10

In a particularly preferred embodiment of the present invention, in a compound of formula I,  $n = 1$ ,  $L_1$  is as defined herein and  $m = 0$ .

15 Preferably, when  $n = 1$  in the compound of formula I,  $L_1$  represents cyclopentadienyl, indenyl or aryl ring each of which rings are optionally substituted by one or more substituents selected from hydrogen, lower alkyl, halo, cyano, nitro,  $-OR^{19}$ ,  $-OC(O)R^{20}$ ,  $-C(O)R^{21}$ ,  $-C(O)OR^{22}$ ,  
20  $-N(R^{23})R^{24}$ ,  $-C(O)N(R^{25})R^{26}$ ,  $-C(S)(R^{27})R^{28}$ ,  $-SR^{29}$ ,  $-C(O)SR^{30}$ ,  
 $-CF_3$  or ferrocenyl (by which we mean the cyclopentadienyl, indenyl or aryl ring which  $L_1$  may represent is bonded directly to the cyclopentadienyl ring of the ferrocenyl group), wherein  $R^{19}$  to  $R^{30}$  is as defined herein. More  
25 preferably, if the cyclopentadienyl, indenyl or aryl ring which  $L_1$  may represent is substituted it is preferably substituted with one or more substituents selected from unsubstituted  $C_1$ - $C_6$  alkyl, halo, cyano,  $-OR^{19}$ ,  $-OC(O)R^{20}$ ,  
 $-C(O)R^{21}$ ,  $-C(O)OR^{22}$ ,  $-N(R^{23})R^{24}$  where  $R^{19}$ ,  $R^{20}$ ,  $R^{21}$ ,  $R^{22}$ ,  $R^{23}$   
30 and  $R^{24}$  each independently represent hydrogen or  $C_1$ - $C_6$  alkyl. Even more preferably, if the cyclopentadienyl, indenyl or aryl ring which  $L_1$  may represent is

substituted, it is preferably substituted with one or more substituents selected from unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl.

Preferably, when  $n = 1$ , L<sub>1</sub> represents cyclopentadienyl, indenyl, phenyl or naphthyl optionally substituted as defined herein. Preferably, the cyclopentadienyl, indenyl, phenyl or naphthyl groups are unsubstituted. More preferably, L<sub>1</sub> represents cyclopentadienyl, indenyl or phenyl, each of which rings are unsubstituted. Most preferably, L<sub>1</sub> represents unsubstituted cyclopentadienyl.

Alternatively, when  $n = 0$ , the compounds of the invention contain only one cyclopentadienyl or indenyl ring (each of which ring systems may optionally be substituted as described herein). Such compounds may be referred to as "half sandwich compounds". Preferably, when  $n = 0$  then  $m$  represents 1 to 5 so that the metal M of the compounds of formula I has an 18 electron count. In other words, when metal M of the compounds of formula I is iron, the total number of electrons contributed by the ligands L<sub>2</sub> is typically five.

In a particularly preferred alternative embodiment of the present invention, in a compound of formula I,  $n = 0$ , L<sub>2</sub> is as defined herein and  $m = 3$  or 4, particularly 3.

Preferably, when  $n$  is equal to zero and  $m$  is not equal to zero in a compound of formula I, L<sub>2</sub> represents one or more ligands each of which are independently selected from lower alkyl, halo, -CO, -P(R<sup>43</sup>)(R<sup>44</sup>)R<sup>45</sup> or -N(R<sup>46</sup>)(R<sup>47</sup>)R<sup>48</sup>. More preferably, L<sub>2</sub> represents one or more ligands each of which are independently selected from unsubstituted C<sub>1</sub> to C<sub>4</sub> alkyl, halo, particularly chloro, -CO, -P(R<sup>43</sup>)(R<sup>44</sup>)R<sup>45</sup> or

$-N(R^{46})(R^{47})R^{48}$ , wherein  $R^{43}$  to  $R^{48}$  are independently selected from hydrogen, unsubstituted  $C_1$  to  $C_6$  alkyl or aryl, such as phenyl.

5 Suitably, the metal M or metal cation thereof in the compounds of formula I is typically bonded to the cyclopentadienyl ring(s), the cyclopentadienyl moiety of the indenyl ring(s) if present, the aryl ring if present, and/or the ligands  $L_2$  if present. Typically, the  
10 cyclopentadienyl ring or the cyclopentadienyl moiety of the indenyl ring exhibits a pentahapto bonding mode with the metal; however other bonding modes between the cyclopentadienyl ring or cyclopentadienyl moiety of the indenyl ring and the metal, such as trihapto coordination,  
15 are also embraced by the scope of the present invention.

Most preferably, in a compound of formula I,  $n = 1$ ,  $m = 0$  and  $L_1$  is defined herein, particularly unsubstituted cyclopentadienyl.

20

By the term "M represents a Group VIB or VIIIB metal" in a compound of formula I we include metals such as Cr, Mo, W, Fe, Co, Ni, Ru and Rh. For the avoidance of doubt, references to Group VIB or VIIIB metals herein include  
25 metals of Groups 6, 8, 9 and 10 in the modern periodic table nomenclature.

By the term "metal cation thereof" we mean that the Group VIB or VIIIB metal (M) in the compound of formula I as  
30 defined herein has a positive charge. Suitably, the metal cation may be in the form of a salt or may comprise weakly coordinated anions derived from halo, nitric acid; sulphuric acid; lower alkanoic (up to  $C_{12}$ ) acids such as

acetic acid and propionic acid; sulphonic acids such as methane sulphonic acid, chlorosulphonic acid, fluorosulphonic acid, trifluoromethane sulphonic acid, benzene sulphonic acid, naphthalene sulphonic acid, 5 toluene sulphonic acid, e.g. p-toluene sulphonic acid, t-butyl sulphonic acid, and 2-hydroxypropane sulphonic acid; sulphonated ion exchange resins; perhalic acid such as perchloric acid; perfluorinated carboxylic acid such as trichloroacetic acid and trifluoroacetic acid; 10 orthophosphoric acid; phosphonic acid such as benzene phosphonic acid; and acids derived from interactions between Lewis acids and Broensted acids. Other sources which may provide suitable anions include the tetraphenyl borate derivatives.

15

Preferably M represents a Group VIB or VIIIB metal. In other words the total electron count for the metal M is 18.

20 Preferably, in the compound of formula I, M represents Cr, Mo, Fe, Co or Ru, or a metal cation thereof. Even more preferably, M represents Cr, Fe, Co or Ru or a metal cation thereof. Most preferably, M is selected from a Group VIIIB metal or metal cation thereof. An especially 25 preferred Group VIIIB metal is Fe. Although the metal M as defined herein may be in a cationic form, preferably it carries essentially no residual charge due to coordination with  $L_1$  and/or  $L_2$  as defined herein.

30 The term "aryl" when used herein, includes six- to ten-membered carbocyclic aromatic groups, such as phenyl and naphthyl, which groups are optionally substituted with one or more substituents selected from aryl, lower alkyl

(which alkyl group may itself be optionally substituted or terminated as defined below), Het, halo, cyano, nitro, -OR<sup>19</sup>, -OC(O)R<sup>20</sup>, -C(O)R<sup>21</sup>, -C(O)OR<sup>22</sup>, -N(R<sup>23</sup>)R<sup>24</sup>, -C(O)N(R<sup>25</sup>)R<sup>26</sup>, -SR<sup>29</sup>, -C(O)SR<sup>30</sup> or -C(S)N(R<sup>27</sup>)R<sup>28</sup> wherein R<sup>19</sup> to R<sup>30</sup> each independently represent hydrogen, aryl or lower alkyl (which alkyl group may itself be optionally substituted or terminated as defined below). Preferred aryl groups which k, D, E, R<sup>1</sup> to R<sup>55</sup> and L<sub>1</sub> may represent and which adamantyl, 2-phospha-adamantyl, congressyl and lower alkyl may be substituted include phenyl which is optionally substituted with one or more substituents selected from C<sub>1</sub>-C<sub>6</sub> alkyl (which alkyl group itself may be optionally substituted or terminated as defined below), OR<sup>19</sup>, R<sup>19</sup>, halo and NR<sup>23</sup>(R<sup>24</sup>), where R<sup>19</sup>, R<sup>23</sup> and R<sup>24</sup> independently represent hydrogen or lower alkyl. Further preferred aryl groups include phenyl which is optionally substituted with one or more substituents selected from unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl, OR<sup>19</sup> wherein R<sup>19</sup> represents hydrogen or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl, particularly unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl.

The term "Het", when used herein, includes four- to twelve-membered, preferably four- to ten-membered ring systems, which rings contain one or more heteroatoms selected from nitrogen, oxygen, sulfur and mixtures thereof, and which rings contain one or more double bonds or be non-aromatic, partly aromatic or wholly aromatic in character. The ring systems may be monocyclic, bicyclic or fused. Each "Het" group identified herein is optionally substituted by one or more substituents selected from halo, cyano, nitro, oxo, lower alkyl (which alkyl group may itself be optionally substituted or terminated as defined below) -OR<sup>19</sup>, -OC(O)R<sup>20</sup>, -C(O)R<sup>21</sup>, -C(O)OR<sup>22</sup>,

$-N(R^{23})R^{24}$ ,  $-C(O)N(R^{25})R^{26}$ ,  $-SR^{29}$ ,  $-C(O)SR^{30}$  or  $-C(S)N(R^{27})R^{28}$  wherein  $R^{19}$  to  $R^{30}$  each independently represent hydrogen, aryl or lower alkyl (which alkyl group itself may be optionally substituted or terminated as defined below).

5 The term "Het" thus includes groups such as optionally substituted azetidiny, pyrrolidinyl, imidazolyl, indolyl, furanyl, oxazolyl, isoxazolyl, oxadiazolyl, thiazolyl, thiadiazolyl, triazolyl, oxatriazolyl, thiatriazolyl, pyridazinyl, morpholinyl, pyrimidinyl, pyrazinyl,  
10 quinolinyl, isoquinolinyl, piperidinyl, pyrazolyl and piperazinyl. Substitution at Het may be at a carbon atom of the Het ring or, where appropriate, at one or more of the heteroatoms.

15 "Het" groups may also be in the form of an N oxide.

The term "lower alkyl" when used herein, means  $C_1$  to  $C_{10}$  alkyl and includes methyl, ethyl, propyl, butyl, pentyl, hexyl and heptyl groups. Unless otherwise specified, alkyl  
20 groups may, when there is a sufficient number of carbon atoms, be linear or branched, be saturated or unsaturated, be cyclic, acyclic or part cyclic/acyclic, and/or be substituted or terminated by one or more substituents selected from halo, cyano, nitro,  $-OR^{19}$ ,  $-OC(O)R^{20}$ ,  
25  $-C(O)R^{21}$ ,  $-C(O)OR^{22}$ ,  $-N(R^{23})R^{24}$ ,  $-C(O)N(R^{25})R^{26}$ ,  $-SR^{29}$ ,  $-C(O)SR^{30}$ ,  $-C(S)N(R^{27})R^{28}$ , aryl or Het, wherein  $R^{19}$  to  $R^{30}$  each independently represent hydrogen, aryl or lower alkyl, and/or be interrupted by one or more oxygen or sulfur atoms, or by silano or dialkylsilicon groups.

30

Lower alkyl groups which  $R^1$  to  $R^{62}$ , K, D, E and  $L_2$  may represent and which aryl, Het and  $L_1$  may be substituted, may, when there is a sufficient number of carbon atoms, be

linear or branched, be saturated or unsaturated, be cyclic, acyclic or part cyclic/acyclic, and/or be interrupted by one or more of oxygen or sulfur atoms, or by silano or dialkylsilicon groups, and/or be substituted  
5 by one or more substituents selected from halo, cyano, nitro,  $-OR^{19}$ ,  $-OC(O)R^{20}$ ,  $-C(O)R^{21}$ ,  $-C(O)OR^{22}$ ,  $-N(R^{23})R^{24}$ ,  $-C(O)N(R^{25})R^{26}$ ,  $-SR^{29}$ ,  $-C(O)SR^{30}$ ,  $-C(S)N(R^{27})R^{28}$ , aryl or Het wherein  $R^{19}$  to  $R^{30}$  each independently represent hydrogen, aryl or lower alkyl.

10

Similarly, the term "lower alkylene" which  $A_1$  and  $A_2$ , and  $A_3$ ,  $A_4$  and  $A_5$  (when present), represent in a compound of formula I, when used herein, includes  $C_1$  to  $C_{10}$  groups which can be bonded at two places on the group and is  
15 otherwise defined in the same way as "lower alkyl".

Halo groups, which  $L_2$  may represent and with which the above-mentioned groups may be substituted or terminated, include fluoro, chloro, bromo and iodo.

20

Where a compound of the formula (I) contains an alkenyl group, cis (E) and trans (Z) isomerism may also occur. The present invention includes the individual stereoisomers of the compounds of formula (I) and, where appropriate, the  
25 individual tautomeric forms thereof, together with mixtures thereof. Separation of diastereoisomers or cis and trans isomers may be achieved by conventional techniques, e.g. by fractional crystallisation, chromatography or H.P.L.C. of a stereoisomeric mixture of  
30 a compound of the formula (I) or a suitable salt or derivative thereof. An individual enantiomer of a compound of the formula (I) may also be prepared from a corresponding optically pure intermediate or by

resolution, such as by H.P.L.C. of the corresponding racemate using a suitable chiral support or by fractional crystallisation of the diastereoisomeric salts formed by reaction of the corresponding racemate with a suitable  
5 optically active acid or base, as appropriate.

All stereoisomers are included within the scope of the invention.

- 10 Suitable Group VIIIB metals or a compound thereof which may be combined with a compound of formula I thereby forming the compounds of the invention include cobalt, nickel, palladium, rhodium and platinum. Preferably, the Group VIIIB metal is palladium or a compound thereof.
- 15 Suitable compounds of such Group VIIIB metals include salts of such metals with, or compounds comprising weakly coordinated anions derived from, nitric acid; sulphuric acid; lower alkanolic (up to C<sub>12</sub>) acids such as acetic acid and propionic acid; sulphonic acids such as methane  
20 sulphonic acid, chlorosulphonic acid, fluorosulphonic acid, trifluoromethane sulphonic acid, benzene sulphonic acid, naphthalene sulphonic acid, toluene sulphonic acid, e.g. p-toluene sulphonic acid, t-butyl sulphonic acid, and 2-hydroxypropane sulphonic acid; sulphonated ion exchange  
25 resins; perhalic acid such as perchloric acid; perfluorinated carboxylic acid such as trichloroacetic acid and trifluoroacetic acid; orthophosphoric acid; phosphonic acid such as benzene phosphonic acid; and acids derived from interactions between Lewis acids and  
30 Brønsted acids. Other sources, which may provide suitable anions, include the tetraphenyl borate derivatives. Additionally, zero valent palladium with labile ligands e.g. tri(dibenzylideneacetone)dipalladium may be used.



Preferably, zero valent Group VIIIIB metals with labile ligands are employed.

Suitably, the compounds of the invention when employed to  
5 catalyse the carbonylation of ethylenically unsaturated  
compounds include a source of anions. Conveniently, the  
source of anions may be derived by combining a Group VIIIIB  
compound thereof as described in the preceding paragraph  
with a compound of formula I. Alternatively, or  
10 additionally, a separate source of anions as mentioned  
above may be added to the compounds of the present  
invention. Preferably, the source of anions are derived  
from an acid having a pKa less than 4, more preferably a  
pKa less than 3, as measured at 18°C in an aqueous  
15 solution.

Especially preferred compounds of formula I include those  
wherein:

- 20 (1)  $X^1$  represents  $CR^1(R^2)(R^3)$ ,  $X^2$  represents  $CR^4(R^5)(R^6)$ ,  
 $X^3$  represents  $CR^7(R^8)(R^9)$ ,  $X^4$  represents  $CR^{10}(R^{11})(R^{12})$ ,  
wherein each of  $R^1$  to  $R^{12}$  independently represents  
unsubstituted  $C_1$ - $C_6$  alkyl or trifluoromethyl,  
particularly where each of  $R^1$  to  $R^{12}$  is identical,  
25 especially where each of  $R^1$  to  $R^{12}$  represents  
unsubstituted  $C_1$ - $C_6$  alkyl, particularly methyl;  
 $A_1$  and  $A_2$  are the same and represent  $-CH_2-$ ;  
 $K$ ,  $D$  and  $E$  are the same and represent hydrogen or  
unsubstituted  $C_1$ - $C_6$  alkyl, particularly hydrogen;  
30  $Q^1$  and  $Q^2$  both represent phosphorus;  
 $M$  represents Fe;

$n = 1$  and  $L_1$  represents cyclopentadienyl, particularly unsubstituted cyclopentadienyl, and  $m = 0$ .

5 (2)  $X^1$  represents  $CR^1(R^2)(R^3)$ ,  $X^2$  represents  $CR^4(R^5)(R^6)$ ,  $X^3$  represents  $CR^7(R^8)(R^9)$ ,  $X^4$  represents  $CR^{10}(R^{11})(R^{12})$ ;  $K$  represents  $-CH_2-Q^3(X^5)X^6$  wherein  $X^5$  represents  $CR^{13}(R^{14})(R^{15})$  and  $X^6$  represents  $CR^{16}(R^{17})(R^{18})$ ; each of  $R^1$  to  $R^{18}$  independently represent  
10 unsubstituted  $C_1-C_6$  alkyl or trifluoromethyl, particularly where each of  $R^1$  to  $R^{18}$  is identical, especially where each of  $R^1$  to  $R^{18}$  represents unsubstituted  $C_1-C_6$  alkyl, particularly methyl;  $A_1$  and  $A_2$  are the same and represent  $-CH_2-$ ;  
15  $Q^1$ ,  $Q^2$  and  $Q^3$  each represent phosphorus;  $D$  and  $E$  are the same and represent hydrogen or unsubstituted  $C_1-C_6$  alkyl, particularly hydrogen;  $M$  represents Fe;  
 $n = 1$  and  $L_1$  represents cyclopentadienyl,  
20 particularly unsubstituted cyclopentadienyl, and  $m = 0$ .

(3)  $X^1$  represents  $CR^1(R^2)(R^3)$ ,  $X^2$  represents  $CR^4(R^5)(R^6)$ ,  $X^3$  represents  $CR^7(R^8)(R^9)$ ,  $X^4$  represents  $CR^{10}(R^{11})(R^{12})$ ;  $K$  represents  $-CH_2-Q^3(X^5)X^6$  wherein  $X^5$  represents  $CR^{13}(R^{14})(R^{15})$  and  $X^6$  represents  $CR^{16}(R^{17})(R^{18})$ ;  
25 each of  $R^1$  to  $R^{18}$  independently represent unsubstituted  $C_1-C_6$  alkyl or trifluoromethyl, particularly where each of  $R^1$  to  $R^{18}$  is identical, especially where each of  $R^1$  to  $R^{18}$  represents  
30 unsubstituted  $C_1-C_6$  alkyl, particularly methyl;  $A_1$  and  $A_2$  are the same and represent  $-CH_2-$ ;  
 $Q^1$ ,  $Q^2$  and  $Q^3$  each represent phosphorus;

D and E together with the carbon atoms of the cyclopentadienyl ring to which they are attached form an unsubstituted phenyl ring;

M represents Fe;

5         $n = 1$  and  $L_1$  represents cyclopentadienyl, particularly unsubstituted cyclopentadienyl, and  $m = 0$ .

(4)     $X^1$  represents  $CR^1(R^2)(R^3)$ ,  $X^2$  represents  $CR^4(R^5)(R^6)$ ,  
 10         $X^3$  represents  $CR^7(R^8)(R^9)$ ,  $X^4$  represents  $CR^{10}(R^{11})(R^{12})$ ,  
 wherein each of  $R^1$  to  $R^{12}$  independently represent unsubstituted  $C_1$ - $C_6$  alkyl or trifluoromethyl, particularly where each of  $R^1$  to  $R^{12}$  is identical, especially where each of  $R^1$  to  $R^{12}$  represents  
 15        unsubstituted  $C_1$ - $C_6$  alkyl, particularly methyl;

$A_1$  and  $A_2$  are the same and represent  $-CH_2-$ ;

$Q^1$  and  $Q^2$  both represent phosphorus;

K represents hydrogen or  $C_1$ - $C_6$  alkyl, particularly hydrogen;

20        D and E together with the carbon atoms of the cyclopentadienyl ring to which they are attached form an unsubstituted phenyl ring;

M represents Fe;

25         $n = 1$  and  $L_1$  represents cyclopentadienyl, particularly unsubstituted cyclopentadienyl, and  $m = 0$ .

(5)     $X^1$  represents  $CR^1(R^2)(R^3)$ ,  $X^2$  represents  $CR^4(R^5)(R^6)$ ,  
 $X^3$  represents  $CR^7(R^8)(R^9)$ ,  $X^4$  represents  $CR^{10}(R^{11})(R^{12})$ ;  
 30        E represents  $-CH_2-Q^5(X^9)X^{10}$  wherein  $X^9$  represents  $CR^{37}(R^{38})(R^{39})$  and  $X^{10}$  represents  $CR^{40}(R^{41})(R^{42})$ ;  
 each of  $R^1$  to  $R^{12}$  and  $R^{37}$  to  $R^{42}$  independently represent unsubstituted  $C_1$ - $C_6$  alkyl or

trifluoromethyl, particularly where each of  $R^1$  to  $R^{12}$  and  $R^{37}$  to  $R^{42}$  is identical, especially where each of  $R^1$  to  $R^{12}$  and  $R^{37}$  to  $R^{42}$  represents unsubstituted  $C_1$ - $C_6$  alkyl, particularly methyl;

5  $A_1$  and  $A_2$  are the same and represent  $-CH_2-$ ;

$Q^1$ ,  $Q^2$  and  $Q^5$  each represent phosphorus;

$D$  and  $K$  are the same and represent hydrogen or unsubstituted  $C_1$ - $C_6$  alkyl, particularly hydrogen;

$M$  represents Fe;

10  $n = 1$  and  $L_1$  represents cyclopentadienyl, particularly unsubstituted cyclopentadienyl, and  $m = 0$ .

(6)  $X^1$  represents  $CR^1(R^2)(R^3)$ ,  $X^2$  represents  $CR^4(R^5)(R^6)$ ,  
15  $X^3$  represents  $CR^7(R^8)(R^9)$ ,  $X^4$  represents  $CR^{10}(R^{11})(R^{12})$ ;  
 $K$  represents  $-CH_2-Q^3(X^5)X^6$  wherein  $X^5$  represents  $CR^{13}(R^{14})(R^{15})$  and  $X^6$  represents  $CR^{16}(R^{17})(R^{18})$ ;

$D$  represents  $-CH_2-Q^4(X^7)X^8$  wherein  $X^7$  represents  $CR^{31}(R^{32})(R^{33})$  and  $X^8$  represents  $CR^{34}(R^{35})(R^{36})$ ;

20  $E$  represents  $-CH_2-Q^5(X^9)X^{10}$  wherein  $X^9$  represents  $CR^{37}(R^{38})(R^{39})$  and  $X^{10}$  represents  $CR^{40}(R^{41})(R^{42})$ ;

each of  $R^1$  to  $R^{18}$  and  $R^{31}$  to  $R^{42}$  independently represent unsubstituted  $C_1$ - $C_6$  alkyl or trifluoromethyl, particularly where each of  $R^1$  to  $R^{18}$   
25 and  $R^{31}$  to  $R^{42}$  is identical, especially where each of  $R^1$  to  $R^{18}$  and  $R^{31}$  to  $R^{42}$  represents unsubstituted  $C_1$ - $C_6$  alkyl, particularly methyl;

$A_1$  and  $A_2$  are the same and represent  $-CH_2-$ ;

$Q^1$ ,  $Q^2$ ,  $Q^3$ ,  $Q^4$  and  $Q^5$  each represent phosphorus

30  $M$  represents Fe;

$n = 1$  and  $L_1$  represents cyclopentadienyl, particularly unsubstituted cyclopentadienyl, and  $m = 0$ .

- (7)  $X^1$ ,  $X^2$ ,  $X^3$  and  $X^4$  independently represent adamantyl, especially where  $X^1$  to  $X^4$  represent the same adamantyl group;
- 5  $A_1$  and  $A_2$  are the same and represent  $-\text{CH}_2-$ ;  
 $K$ ,  $D$  and  $E$  are the same and represent hydrogen or unsubstituted  $\text{C}_1$ - $\text{C}_6$  alkyl, particularly hydrogen;  
 $Q^1$  and  $Q^2$  both represent phosphorus;  
 $M$  represents Fe;
- 10  $n = 1$  and  $L_1$  represents cyclopentadienyl, particularly unsubstituted cyclopentadienyl, and  $m = 0$ .
- (8)  $X^1$ ,  $X^2$ ,  $X^3$  and  $X^4$  independently represent adamantyl, especially where  $X^1$  to  $X^4$  represent the same adamantyl group;
- 15  $K$  represents  $-\text{CH}_2-\text{Q}^3(\text{X}^5)\text{X}^6$  wherein  $\text{X}^5$  and  $\text{X}^6$  independently represent adamantyl, especially where  $\text{X}^1$  to  $\text{X}^6$  represent the same adamantyl group;
- 20  $A_1$  and  $A_2$  are the same and represent  $-\text{CH}_2-$ ;  
 $Q^1$ ,  $Q^2$  and  $Q^3$  each represent phosphorus;  
 $D$  and  $E$  are the same and represent hydrogen or unsubstituted  $\text{C}_1$ - $\text{C}_6$  alkyl, particularly hydrogen;  
 $M$  represents Fe;
- 25  $n = 1$  and  $L_1$  represents cyclopentadienyl, particularly unsubstituted cyclopentadienyl, and  $m = 0$ .
- (9)  $X^1$ ,  $X^2$ ,  $X^3$  and  $X^4$  independently represent adamantyl, especially where  $X^1$  to  $X^4$  represent the same adamantyl group;
- 30

- K represents  $-\text{CH}_2-\text{Q}^3(\text{X}^5)\text{X}^6$  wherein  $\text{X}^5$  and  $\text{X}^6$  independently represent adamantyl, especially where  $\text{X}^1$  to  $\text{X}^6$  represent the same adamantyl group;  
 $\text{A}_1$  and  $\text{A}_2$  are the same and represent  $-\text{CH}_2-$ ;  
 5  $\text{Q}^1$ ,  $\text{Q}^2$  and  $\text{Q}^3$  each represent phosphorus;  
 D and E together with the carbon atoms of the cyclopentadienyl ring to which they are attached form an unsubstituted phenyl ring;  
 M represents Fe;  
 10  $n = 1$  and  $\text{L}_1$  represents cyclopentadienyl, particularly unsubstituted cyclopentadienyl, and  $m = 0$ .
- (10)  $\text{X}^1$ ,  $\text{X}^2$ ,  $\text{X}^3$  and  $\text{X}^4$  independently represent adamantyl, especially where  $\text{X}^1$  to  $\text{X}^4$  represent the same adamantyl group;  
 15  $\text{A}_1$  and  $\text{A}_2$  are the same and represent  $-\text{CH}_2-$ ;  
 $\text{Q}^1$  and  $\text{Q}^2$  both represent phosphorus;  
 K represents hydrogen or unsubstituted  $\text{C}_1$ - $\text{C}_6$  alkyl, particularly hydrogen;  
 20 D and E together with the carbon atoms of the cyclopentadienyl ring to which they are attached form an unsubstituted phenyl ring;  
 M represents Fe;  
 25  $n = 1$  and  $\text{L}_1$  represents cyclopentadienyl, particularly unsubstituted cyclopentadienyl, and  $m = 0$ .
- (11)  $\text{X}^1$ ,  $\text{X}^2$ ,  $\text{X}^3$  and  $\text{X}^4$  independently represent adamantyl;  
 30 K represents  $-\text{CH}_2-\text{Q}^3(\text{X}^5)\text{X}^6$  wherein  $\text{X}^5$  and  $\text{X}^6$  independently represent adamantyl;  
 D represents  $-\text{CH}_2-\text{Q}^4(\text{X}^7)\text{X}^8$  wherein  $\text{X}^7$  and  $\text{X}^8$  independently represents adamantyl;

E represents  $-\text{CH}_2-\text{Q}^5(\text{X}^9)\text{X}^{10}$  wherein  $\text{X}^9$  and  $\text{X}^{10}$  independently represents adamantyl, especially where  $\text{X}^1$  to  $\text{X}^{10}$  represent the same adamantyl group;

$\text{A}_1$  and  $\text{A}_2$  are the same and represent  $-\text{CH}_2-$ ;

5  $\text{Q}^1$ ,  $\text{Q}^2$ ,  $\text{Q}^3$ ,  $\text{Q}^4$  and  $\text{Q}^5$  each represent phosphorus;

M represents Fe;

$n = 1$  and  $\text{L}_1$  represents cyclopentadienyl, particularly unsubstituted cyclopentadienyl, and  $m = 0$ .

10

(12)  $\text{X}^1$  and  $\text{X}^2$  together with  $\text{Q}^2$  to which they are attached represents 2-phospha-adamantyl;

$\text{X}^3$  and  $\text{X}^4$  together with  $\text{Q}^1$  to which they are attached represents 2-phospha-adamantyl;

15  $\text{A}_1$  and  $\text{A}_2$  are the same and represent  $-\text{CH}_2-$ ;

K, D and E are the same and represent hydrogen or unsubstituted  $\text{C}_1$ - $\text{C}_6$  alkyl, particularly hydrogen;

$\text{Q}^1$  and  $\text{Q}^2$  both represent phosphorus;

M represents Fe;

20

$n = 1$  and  $\text{L}_1$  represents cyclopentadienyl, particularly unsubstituted cyclopentadienyl, and  $m = 0$ .

(13)  $\text{X}^1$  and  $\text{X}^2$  together with  $\text{Q}^2$  to which they are attached represents 2-phospha-adamantyl;

25

$\text{X}^3$  and  $\text{X}^4$  together with  $\text{Q}^1$  to which they are attached represents 2-phospha-adamantyl;

K represents  $-\text{CH}_2-\text{Q}^3(\text{X}^5)\text{X}^6$  wherein  $\text{X}^5$  and  $\text{X}^6$  together with  $\text{Q}^3$  to which they are attached represents 2-phospha-adamantyl;

30

$\text{A}_1$  and  $\text{A}_2$  are the same and represent  $-\text{CH}_2-$ ;

$\text{Q}^1$ ,  $\text{Q}^2$  and  $\text{Q}^3$  each represent phosphorus;

D and E are the same and represent hydrogen or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl, particularly hydrogen;

M represents Fe;

5        n = 1 and L<sub>1</sub> represents cyclopentadienyl, particularly unsubstituted cyclopentadienyl, and m = 0.

(14) X<sup>1</sup> and X<sup>2</sup> together with Q<sup>2</sup> to which they are attached represents 2-phospha-adamantyl;

10        X<sup>3</sup> and X<sup>4</sup> together with Q<sup>1</sup> to which they are attached represents 2-phospha-adamantyl;

K represents -CH<sub>2</sub>-Q<sup>3</sup>(X<sup>5</sup>)X<sup>6</sup> wherein X<sup>5</sup> and X<sup>6</sup> together with Q<sup>3</sup> to which they are attached represents 2-phospha-adamantyl;

15        A<sub>1</sub> and A<sub>2</sub> are the same and represent -CH<sub>2</sub>-;

Q<sup>1</sup>, Q<sup>2</sup> and Q<sup>3</sup> each represent phosphorus;

D and E together with the carbon atoms of the cyclopentadienyl ring to which they are attached form an unsubstituted phenyl ring;

20        M represents Fe;

n = 1 and L<sub>1</sub> represents cyclopentadienyl, particularly unsubstituted cyclopentadienyl, and m = 0.

25        (15) X<sup>1</sup> and X<sup>2</sup> together with Q<sup>2</sup> to which they are attached represents 2-phospha-adamantyl;

X<sup>3</sup> and X<sup>4</sup> together with Q<sup>1</sup> to which they are attached represents 2-phospha-adamantyl;

A<sub>1</sub> and A<sub>2</sub> are the same and represent -CH<sub>2</sub>-;

30        Q<sup>1</sup> and Q<sup>2</sup> both represent phosphorus;

K represents hydrogen or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl, particularly hydrogen;



D and E together with the carbon atoms of the cyclopentadienyl ring to which they are attached form an unsubstituted phenyl ring;

M represents Fe;

5        n = 1 and L<sub>1</sub> represents cyclopentadienyl, particularly unsubstituted cyclopentadienyl, and m = 0.

10        (16) X<sup>1</sup> and X<sup>2</sup> together with Q<sup>2</sup> to which they are attached represents 2-phospha-adamantyl;

X<sup>3</sup> and X<sup>4</sup> together with Q<sup>1</sup> to which they are attached represents 2-phospha-adamantyl;

15        K represents -CH<sub>2</sub>-Q<sup>3</sup>(X<sup>5</sup>)X<sup>6</sup> wherein X<sup>5</sup> and X<sup>6</sup> together with Q<sup>3</sup> to which they are attached represents 2-phospha-adamantyl;

D represents -CH<sub>2</sub>-Q<sup>4</sup>(X<sup>7</sup>)X<sup>8</sup> wherein X<sup>7</sup> and X<sup>8</sup> together with Q<sup>4</sup> to which they are attached represents 2-phospha-adamantyl;

20        E represents -CH<sub>2</sub>-Q<sup>5</sup>(X<sup>9</sup>)X<sup>10</sup> wherein X<sup>9</sup> and X<sup>10</sup> together with Q<sup>5</sup> to which they are attached represents 2-phospha-adamantyl;

A<sub>1</sub> and A<sub>2</sub> are the same and represent -CH<sub>2</sub>-;

Q<sup>1</sup>, Q<sup>2</sup>, Q<sup>3</sup>, Q<sup>4</sup> and Q<sup>5</sup> each represent phosphorus

M represents Fe;

25        n = 1 and L<sub>1</sub> represents cyclopentadienyl, particularly unsubstituted cyclopentadienyl, and m = 0.

30        (17) X<sup>1</sup> and X<sup>2</sup> together with Q<sup>2</sup> to which they are attached form a ring system of formula Ia, X<sup>3</sup> and X<sup>4</sup> together with Q<sup>1</sup> to which they are attached form a ring system of formula Ib, wherein Y<sup>1</sup> and Y<sup>2</sup> both represent oxygen, R<sup>50</sup> to R<sup>53</sup> are independently

- selected from unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl or CF<sub>3</sub>, and R<sup>49</sup> and R<sup>54</sup> represent hydrogen;  
A<sub>1</sub> and A<sub>2</sub> are the same and represent -CH<sub>2</sub>-;  
K, D and E are the same and represent hydrogen or  
5 unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl, particularly hydrogen;  
Q<sup>1</sup> and Q<sup>2</sup> both represent phosphorus;  
M represents Fe;  
n = 1 and L<sub>1</sub> represents cyclopentadienyl,  
particularly unsubstituted cyclopentadienyl  
10 (referred to as puc) and m = 0.
- (18) X<sup>1</sup> and X<sup>2</sup> together with Q<sup>2</sup> to which they are attached  
form a ring system of formula Ia, X<sup>3</sup> and X<sup>4</sup> together  
with Q<sup>1</sup> to which they are attached form a ring  
15 system of formula Ib, wherein Y<sup>1</sup> and Y<sup>2</sup> both  
represent oxygen, R<sup>50</sup> to R<sup>53</sup> are independently  
selected from unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl or CF<sub>3</sub>, and  
R<sup>49</sup> and R<sup>54</sup> represent hydrogen;  
K represents -CH<sub>2</sub>-Q<sup>3</sup>(X<sup>5</sup>)X<sup>6</sup> wherein X<sup>5</sup> and X<sup>6</sup> together  
20 with Q<sup>3</sup> to which they are attached form a ring  
system of formula Ic, wherein Y<sup>3</sup> represents oxygen,  
R<sup>50</sup> to R<sup>53</sup> are independently selected from hydrogen,  
unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl or CF<sub>3</sub> and R<sup>49</sup> and R<sup>54</sup>  
represent hydrogen;  
25 A<sub>1</sub> and A<sub>2</sub> are the same and represent -CH<sub>2</sub>-;  
Q<sup>1</sup>, Q<sup>2</sup> and Q<sup>3</sup> each represent phosphorus;  
D and E are the same and represent hydrogen or C<sub>1</sub>-C<sub>6</sub>  
alkyl, particularly hydrogen;  
M represents Fe;  
30 n = 1 and L<sub>1</sub> represents cyclopentadienyl,  
particularly unsubstituted cyclopentadienyl, and m =  
0.

- (19)  $X^1$  and  $X^2$  together with  $Q^2$  to which they are attached form a ring system of formula Ia,  $X^3$  and  $X^4$  together with  $Q^1$  to which they are attached form a ring system of formula Ib, wherein  $Y^1$  and  $Y^2$  both represent oxygen,  $R^{50}$  to  $R^{53}$  are independently selected from unsubstituted  $C_1$ - $C_6$  alkyl or  $CF_3$ , and  $R^{49}$  and  $R^{54}$  represent hydrogen;  
K represents  $-CH_2-Q^3(X^5)X^6$  wherein  $X^5$  and  $X^6$  together with  $Q^3$  to which they are attached form a ring system of formula Ic, wherein  $Y^3$  represents oxygen,  $R^{50}$  to  $R^{53}$  are independently selected from unsubstituted  $C_1$ - $C_6$  alkyl or  $CF_3$ , and  $R^{49}$  and  $R^{54}$  represent hydrogen;  
 $A_1$  and  $A_2$  are the same and represent  $-CH_2-$ ;  
 $Q^1$ ,  $Q^2$  and  $Q^3$  each represent phosphorus;  
D and E together with the carbon atoms of the cyclopentadienyl ring to which they are attached form an unsubstituted phenyl ring;  
M represents Fe;  
 $n = 1$  and  $L_1$  represents cyclopentadienyl, particularly unsubstituted cyclopentadienyl, and  $m = 0$ .
- (20)  $X^1$  and  $X^2$  together with  $Q^2$  to which they are attached form a ring system of formula Ia,  $X^3$  and  $X^4$  together with  $Q^1$  to which they are attached form a ring system of formula Ib, wherein  $Y^1$  and  $Y^2$  both represent oxygen,  $R^{50}$  to  $R^{53}$  are independently selected from unsubstituted  $C_1$ - $C_6$  alkyl or  $CF_3$ , and  $R^{49}$  and  $R^{54}$  represent hydrogen;  
 $A_1$  and  $A_2$  are the same and represent  $-CH_2-$ ;  
 $Q^1$  and  $Q^2$  both represent phosphorus;

K represents hydrogen or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl, particularly hydrogen;

D and E together with the carbon atoms of the cyclopentadienyl ring to which they are attached form an unsubstituted phenyl ring;

M represents Fe;

n = 1 and L<sub>1</sub> represents cyclopentadienyl, particularly unsubstituted cyclopentadienyl, and m = 0.

10

(21) X<sup>1</sup> and X<sup>2</sup> together with Q<sup>2</sup> to which they are attached form a ring system of formula Ia, X<sup>3</sup> and X<sup>4</sup> together with Q<sup>1</sup> to which they are attached form a ring system of formula Ib, wherein Y<sup>1</sup> and Y<sup>2</sup> both represent oxygen, R<sup>50</sup> to R<sup>53</sup> are independently selected from unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl or CF<sub>3</sub>, and R<sup>49</sup> and R<sup>54</sup> represent hydrogen;

15

K represents -CH<sub>2</sub>-Q<sup>3</sup>(X<sup>5</sup>)X<sup>6</sup> wherein X<sup>5</sup> and X<sup>6</sup> together with Q<sup>3</sup> to which they are attached form a ring system of formula Ic, wherein Y<sup>3</sup> represents oxygen, R<sup>50</sup> to R<sup>53</sup> are independently selected from unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl or CF<sub>3</sub>, and R<sup>49</sup> and R<sup>54</sup> represent hydrogen;

20

D represents -CH<sub>2</sub>-Q<sup>4</sup>(X<sup>7</sup>)X<sup>8</sup> wherein X<sup>7</sup> and X<sup>8</sup> together with Q<sup>4</sup> to which they are attached form a ring system of formula Ic, wherein Y<sup>3</sup> represents oxygen, R<sup>50</sup> to R<sup>53</sup> are independently selected from unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl or CF<sub>3</sub>, and R<sup>49</sup> and R<sup>54</sup> represent hydrogen;

25

E represents -CH<sub>2</sub>-Q<sup>5</sup>(X<sup>9</sup>)X<sup>10</sup> wherein X<sup>9</sup> and X<sup>10</sup> together with Q<sup>5</sup> to which they are attached form a ring system of formula Ie, wherein Y<sup>5</sup> represents oxygen, and R<sup>50</sup> to R<sup>53</sup> are independently selected from

30

unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl or CF<sub>3</sub>, and R<sup>49</sup> and R<sup>54</sup> represent hydrogen;

A<sub>1</sub> and A<sub>2</sub> are the same and represent -CH<sub>2</sub>-;

Q<sup>1</sup>, Q<sup>2</sup>, Q<sup>3</sup>, Q<sup>4</sup> and Q<sup>5</sup> each represent phosphorus;

5 M represents Fe;

n = 1 and L<sub>1</sub> represents cyclopentadienyl; particularly unsubstituted cyclopentadienyl, and m = 0.

10 (22) X<sup>1</sup>, X<sup>2</sup>, X<sup>3</sup> and X<sup>4</sup> independently represent congressyl, especially where X<sup>1</sup> to X<sup>4</sup> represent the same congressyl group;

A<sub>1</sub> and A<sub>2</sub> are the same and represent -CH<sub>2</sub>-;

15 K, D and E are the same and represent hydrogen or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl, particularly hydrogen;

Q<sup>1</sup> and Q<sup>2</sup> both represent phosphorus;

M represents Fe;

20 n = 1 and L<sub>1</sub> represents cyclopentadienyl, particularly unsubstituted cyclopentadienyl, and m = 0.

(23) X<sup>1</sup>, X<sup>2</sup>, X<sup>3</sup> and X<sup>4</sup> independently represent congressyl, especially where X<sup>1</sup> to X<sup>4</sup> represent the same congressyl group;

25 K represents -CH<sub>2</sub>-Q<sup>3</sup>(X<sup>5</sup>)X<sup>6</sup> wherein X<sup>5</sup> and X<sup>6</sup> independently represent congressyl, especially where X<sup>1</sup> to X<sup>6</sup> represent the same congressyl group;

A<sub>1</sub> and A<sub>2</sub> are the same and represent -CH<sub>2</sub>-;

Q<sup>1</sup>, Q<sup>2</sup> and Q<sup>3</sup> each represent phosphorus;

30 D and E are the same and represent hydrogen or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl, particularly hydrogen;

M represents Fe;

$n = 1$  and  $L_1$  represents cyclopentadienyl, particularly unsubstituted cyclopentadienyl, and  $m = 0$ .

5 (24)  $X^1$ ,  $X^2$ ,  $X^3$  and  $X^4$  independently represent congressyl, especially where  $X^1$  to  $X^4$  represent the same congressyl group;

K represents  $-\text{CH}_2-\text{Q}^3(\text{X}^5)\text{X}^6$  wherein  $\text{X}^5$  and  $\text{X}^6$  independently represent congressyl, especially where  
10  $\text{X}^1$  to  $\text{X}^6$  represent the same congressyl group;

$A_1$  and  $A_2$  are the same and represent  $-\text{CH}_2-$ ;

$\text{Q}^1$ ,  $\text{Q}^2$  and  $\text{Q}^3$  each represent phosphorus;

D and E together with the carbon atoms of the cyclopentadienyl ring to which they are attached  
15 form an unsubstituted phenyl ring;

M represents Fe;

$n = 1$  and  $L_1$  represents cyclopentadienyl, particularly unsubstituted cyclopentadienyl, and  $m = 0$ .

20

(25)  $X^1$ ,  $X^2$ ,  $X^3$  and  $X^4$  independently represent congressyl, especially where  $X^1$  to  $X^4$  represent the same congressyl group;

$A_1$  and  $A_2$  are the same and represent  $-\text{CH}_2-$ ;

25  $\text{Q}^1$  and  $\text{Q}^2$  both represent phosphorus;

K represents hydrogen or unsubstituted  $\text{C}_1\text{-C}_6$  alkyl, particularly hydrogen;

D and E together with the carbon atoms of the cyclopentadienyl ring to which they are attached  
30 form an unsubstituted phenyl ring;

M represents Fe;

$n = 1$  and  $L_1$  represents cyclopentadienyl, particularly unsubstituted cyclopentadienyl, and  $m = 0$ .

- 5 (26)  $X^1, X^2, X^3$  and  $X^4$  independently represent congressyl;  
 $K$  represents  $-\text{CH}_2-\text{Q}^3(X^5)X^6$  wherein  $X^5$  and  $X^6$   
independently represent congressyl;  
 $D$  represents  $-\text{CH}_2-\text{Q}^4(X^7)X^8$  wherein  $X^7$  and  $X^8$   
independently represents congressyl;  
10  $E$  represents  $-\text{CH}_2-\text{Q}^5(X^9)X^{10}$  wherein  $X^9$  and  $X^{10}$   
independently represents congressyl, especially  
where  $X^1$  to  $X^{10}$  represent the same congressyl group;  
 $A_1$  and  $A_2$  are the same and represent  $-\text{CH}_2-$ ;  
 $\text{Q}^1, \text{Q}^2, \text{Q}^3, \text{Q}^4$  and  $\text{Q}^5$  each represent phosphorus;  
15  $M$  represents Fe;  
 $n = 1$  and  $L_1$  represents cyclopentadienyl,  
particularly unsubstituted cyclopentadienyl, and  $m = 0$ .
- 20 (27)  $X^1$  and  $X^3$  independently represent adamantyl,  
especially where  $X^1$  and  $X^3$  represent the same  
adamantyl group;  
 $X^2$  represents  $\text{CR}^4(\text{R}^5)(\text{R}^6)$  and  $X^4$  represents  
 $\text{CR}^{10}(\text{R}^{11})(\text{R}^{12})$  wherein each of  $\text{R}^4, \text{R}^5, \text{R}^6, \text{R}^{10}, \text{R}^{11}$  and  
25  $\text{R}^{12}$  independently represent  $\text{C}_1\text{-C}_6$  alkyl or  
trifluoromethyl, particularly where each of  $\text{R}^4$  to  $\text{R}^6$   
and  $\text{R}^{10}$  to  $\text{R}^{12}$  is identical, especially where each of  
 $\text{R}^4$  to  $\text{R}^6$  and  $\text{R}^{10}$  to  $\text{R}^{12}$  represents unsubstituted  $\text{C}_1\text{-C}_6$   
alkyl, particularly methyl;  
30  $A_1$  and  $A_2$  are the same and represent  $-\text{CH}_2-$ ;  
 $K, D$  and  $E$  are the same and represent hydrogen or  
unsubstituted  $\text{C}_1\text{-C}_6$  alkyl, particularly hydrogen;  
 $\text{Q}^1$  and  $\text{Q}^2$  both represent phosphorus;

M represents Fe;

$n = 1$  and  $L_1$  represents cyclopentadienyl, particularly unsubstituted cyclopentadienyl, and  $m = 0$ .

5

(28)  $X^1$  and  $X^3$  independently represent adamantyl, especially where  $X^1$  and  $X^3$  represent the same adamantyl group;

10 K represents  $-\text{CH}_2-\text{Q}^3(\text{X}^5)\text{X}^6$  wherein  $\text{X}^5$  represents adamantyl, especially where  $X^1$ ,  $X^3$  and  $\text{X}^5$  represent the same adamantyl group;

15  $\text{X}^2$  represents  $\text{CR}^4(\text{R}^5)(\text{R}^6)$ ,  $\text{X}^4$  represents  $\text{CR}^{10}(\text{R}^{11})(\text{R}^{12})$ ,  $\text{X}^6$  represents  $\text{CR}^{16}(\text{R}^{17})(\text{R}^{18})$ , wherein each of  $\text{R}^4$  to  $\text{R}^6$ ,  $\text{R}^{10}$  to  $\text{R}^{12}$  and  $\text{R}^{16}$  to  $\text{R}^{18}$  independently represent unsubstituted  $\text{C}_1\text{-C}_6$  alkyl or trifluoromethyl, particularly where each of  $\text{R}^4$  to  $\text{R}^6$ ,  $\text{R}^{10}$  to  $\text{R}^{12}$ , and  $\text{R}^{16}$  to  $\text{R}^{18}$  is identical, especially where each of  $\text{R}^4$  to  $\text{R}^6$ ,  $\text{R}^{10}$  to  $\text{R}^{12}$  and  $\text{R}^{16}$  to  $\text{R}^{18}$  represents unsubstituted  $\text{C}_1\text{-C}_6$  alkyl, particularly methyl;

20  $\text{A}_1$  and  $\text{A}_2$  are the same and represent  $-\text{CH}_2-$ ;

$\text{Q}^1$ ,  $\text{Q}^2$  and  $\text{Q}^3$  each represent phosphorus;

D and E are the same and represent hydrogen or unsubstituted  $\text{C}_1\text{-C}_6$  alkyl, particularly hydrogen;

M represents Fe;

25  $n = 1$  and  $L_1$  represents cyclopentadienyl, particularly unsubstituted cyclopentadienyl, and  $m = 0$ .

30 (29)  $X^1$  and  $X^3$  independently represent adamantyl, especially where  $X^1$  and  $X^3$  represent the same adamantyl group;



- K represents  $-\text{CH}_2-\text{Q}^3(\text{X}^5)\text{X}^6$  wherein  $\text{X}^5$  represents adamantyl, especially where  $\text{X}^1$ ,  $\text{X}^3$  and  $\text{X}^5$  represent the same adamantyl group;
- $\text{X}^2$  represents  $\text{CR}^4(\text{R}^5)(\text{R}^6)$ ,  $\text{X}^4$  represents  $\text{CR}^{10}(\text{R}^{11})(\text{R}^{12})$ ,  
 5  $\text{X}^6$  represents  $\text{CR}^{16}(\text{R}^{17})(\text{R}^{18})$ , wherein each of  $\text{R}^4$  to  $\text{R}^6$ ,  $\text{R}^{10}$  to  $\text{R}^{12}$  and  $\text{R}^{16}$  to  $\text{R}^{18}$  independently represent unsubstituted  $\text{C}_1\text{-C}_6$  alkyl or trifluoromethyl, particularly where each of  $\text{R}^4$  to  $\text{R}^6$ ,  $\text{R}^{10}$  to  $\text{R}^{12}$ , and  $\text{R}^{16}$  to  $\text{R}^{18}$  is identical, especially where each of  $\text{R}^4$   
 10 to  $\text{R}^6$ ,  $\text{R}^{10}$  to  $\text{R}^{12}$  and  $\text{R}^{16}$  to  $\text{R}^{18}$  represents unsubstituted  $\text{C}_1\text{-C}_6$  alkyl, particularly methyl;
- $\text{A}_1$  and  $\text{A}_2$  are the same and represent  $-\text{CH}_2-$ ;
- $\text{Q}^1$ ,  $\text{Q}^2$  and  $\text{Q}^3$  each represent phosphorus;
- D and E together with the carbon atoms of the  
 15 cyclopentadienyl ring to which they are attached form an unsubstituted phenyl ring;
- M represents Fe;
- $n = 1$  and  $\text{L}_1$  represents cyclopentadienyl, particularly unsubstituted cyclopentadienyl, and  $m =$   
 20 0.
- (30)  $\text{X}^1$  and  $\text{X}^3$  independently represent adamantyl, especially where  $\text{X}^1$  and  $\text{X}^3$  represent the same adamantyl group;
- $\text{X}^2$  represents  $\text{CR}^4(\text{R}^5)(\text{R}^6)$  and  $\text{X}^4$  represents  $\text{CR}^{10}(\text{R}^{11})(\text{R}^{12})$  wherein each of  $\text{R}^4$ ,  $\text{R}^5$ ,  $\text{R}^6$ ,  $\text{R}^{10}$ ,  $\text{R}^{11}$  and  $\text{R}^{12}$  independently represent  $\text{C}_1\text{-C}_6$  alkyl or trifluoromethyl, particularly where each of  $\text{R}^4$  to  $\text{R}^6$  and  $\text{R}^{10}$  to  $\text{R}^{12}$  is identical, especially where each of  
 25  $\text{R}^4$  to  $\text{R}^6$  and  $\text{R}^{10}$  to  $\text{R}^{12}$  represents unsubstituted  $\text{C}_1\text{-C}_6$  alkyl, particularly methyl;
- $\text{A}_1$  and  $\text{A}_2$  are the same and represent  $-\text{CH}_2-$ ;
- $\text{Q}^1$  and  $\text{Q}^2$  both represent phosphorus;
- 30

K represents hydrogen or unsubstituted C<sub>1</sub>-C<sub>6</sub> alkyl, particularly hydrogen;

D and E together with the carbon atoms of the cyclopentadienyl ring to which they are attached form an unsubstituted phenyl ring;

M represents Fe;

n = 1 and L<sub>1</sub> represents cyclopentadienyl, particularly unsubstituted cyclopentadienyl, and m = 0.

According to a second aspect, the present invention provides a process for preparing the compounds of the invention comprising combining (a) a Group VIIIB metal or compound thereof, as defined herein; with (b) a compound of formula I as defined herein.

Conveniently, the compounds of the invention may be obtained by dissolving the Group VIIIB metal or compound thereof as defined herein in a suitable solvent such as the ultimate end product of the carbonylation reaction, for example methylpropanoate where the ethylenically unsaturated compound to be carbonylated is ethene in the presence of methanol or methylnonanoate where it is intended to carbonylate octene in the presence of methanol. Preferably, the reactants are mixed at room temperature under an inert atmosphere (e.g. under nitrogen). The molar ratio of the compound of formula I (referred to as (b)) to the Group VIIIB metal or compound thereof (referred to as (a)) is preferably in the range of 1:1 to 5:1, more preferably in the range of 1:1 to 3:1, most preferably in the range of 1:1 to 1:1.25. Conveniently, the possibility of applying these low molar ratios is advantageous, as it avoids the use of an excess


of the compound of formula I and hence minimises the consumption of these usually expensive compounds. Suitably, the compounds of the invention are prepared in a separate step preceding their use in-situ in the carbonylation reaction of an ethylenically unsaturated compound. As mentioned previously, the compounds of the invention may include additional anions derivable from the Group VIIIIB compound thereof, if one is employed, and/or by the addition of a separate source of anions. If a separate source of anions is employed these may be added to the compounds of the invention prior to use in the carbonylation reaction. Alternatively, or additionally, a separate source of anions may be added to the carbonylation reaction.

15



According to a third aspect, the present invention provides a compound of formula I as defined herein.

According to a fourth aspect, the present invention provides a process for the preparation of a compound of formula I, which comprises the reaction of a compound of formula II wherein  $A_1$ ,  $A_2$ ,  $K$ ,  $D$ ,  $E$ ,  $M$ ,  $L_1$ ,  $L_2$ ,  $n$  and  $m$  are as defined for a compound of formula I, and  $LG_1$  and  $LG_2$  represent suitable leaving groups,

25



with a compound of formula IIIa and IIIb



wherein  $X^1$  represents  $CR^1(R^2)(R^3)$ , congressyl or adamantyl,  
 $X^2$  represents  $CR^4(R^5)(R^6)$ , congressyl or adamantyl, or  $X^1$   
 and  $X^2$  together with  $Q^2$  to which they are attached form a  
 10 2-phospha-adamantyl group, or  $X^1$  and  $X^2$  together with  $Q^2$  to  
 which they are attached form a ring system of formula Ia;

wherein  $X^3$  represents  $CR^7(R^8)(R^9)$ , congressyl or adamantyl,  
 $X^4$  represents  $CR^{10}(R^{11})(R^{12})$ , congressyl or adamantyl, or  $X^3$   
 15 and  $X^4$  together with  $Q^1$  to which they are attached form a  
 2-phospha-adamantyl group, or  $X^3$  and  $X^4$  together with  $Q^1$  to  
 which they are attached form a ring system of formula Ib;

$R^1$  to  $R^{12}$ , adamantyl, congressyl,  $Q^1$  and  $Q^2$ , 2-phospha-  
 20 adamantyl, and the ring systems of formula Ia and Ib are  
 as defined for a compound of formula I.

Suitable leaving groups which  $LG_1$  and  $LG_2$  may  
 independently represent include groups which are readily  
 25 displaced by nucleophilic attack by the phosphine, arsine  
 or stibene derivatives IIIa and IIIb. Examples of such  
 groups include halo, particularly bromo and iodo,  $-NR^{23}R^{24}$   
 where  $R^{23}$  and  $R^{24}$  both represent lower alkyl, particularly  
 methyl, and hydroxyl whether in a protonated form or not.  
 30 Preferably,  $LG_1$  and  $LG_2$  each independently represent  $-NMe_2$   
 or hydroxyl. Most preferably, both  $LG_1$  and  $LG_2$  represent  $-$   
 $NMe_2$ , or  $LG_1$  and  $LG_2$  represent  $NMe_2$  or hydroxyl,  
 respectively.

The reaction may be accomplished using methods which are well known to those skilled in the art. For example, the reaction may be accomplished by heating a solution of a compound of formula II with a compound of formula IIIa and  
5 IIIb in anhydrous acetic acid at a temperature between 70 to 90°C, preferably approximately 80°C under an inert atmosphere, such as a nitrogen atmosphere.

10 Preferably, when the compound of formula IIIa represents  $HQ^2(CR^1(R^2)(R^3))CR^4(R^5)(R^6)$  then the compound of formula IIIb represents  $HQ^1(CR^7(R^8)(R^9))CR^{10}(R^{11})(R^{12})$ , wherein  $Q^1$  is the same as  $Q^2$  and  $R^1$  is the same as  $R^7$ ,  $R^2$  is the same as  $R^8$ ,  $R^3$  is the same as  $R^9$ ,  $R^4$  is the same as  $R^{10}$ ,  $R^5$  is the  
15 same as  $R^{11}$ , and  $R^6$  is the same as  $R^{12}$ .

Preferably, when the compound of formula IIIa represents  $HQ^2(X^1)X^2$  wherein  $X^1$  and  $X^2$  independently represent adamantyl, then the compound of formula IIIb represents  
20  $HQ^1(X^3)X^4$  wherein  $X^3$  and  $X^4$  independently represent adamantyl. More preferably,  $X^1$  represents the same adamantyl group as  $X^3$ ,  $X^2$  represents the same adamantyl group as  $X^4$ , and  $Q^1$  is the same as  $Q^2$ .

25 Preferably, when the compound of formula IIIa represents  $HQ^2(X^1)X^2$  wherein  $X^1$  and  $X^2$  independently represent congressyl, then the compound of formula IIIb represents  $HQ^1(X^3)X^4$  wherein  $X^3$  and  $X^4$  independently represent congressyl. More preferably,  $X^1$  represents the same  
30 congressyl group as  $X^3$ ,  $X^2$  represents the same congressyl group as  $X^4$ , and  $Q^1$  is the same as  $Q^2$ .

Preferably, when the compound of formula IIIa represents  $HQ^2(X^1)X^2$  wherein  $X^1$  and  $X^2$  together with  $Q^2$  to which they are attached form a 2-phospha-adamantyl group, then the compound of formula IIIb represents  $HQ^1(X^3)X^4$  wherein  $X^3$  and  $X^4$  together with  $Q^1$  to which they are attached form a 2-phospha-adamantyl group. More preferably,  $X^1$  and  $X^2$  together with  $Q^2$  forms the same 2-phospha-adamantyl group formed by the combination of  $X^3$  and  $X^4$  together with  $Q^1$ .

10 Preferably, when the compound of formula IIIa represents  $HQ^2(X^1)X^2$  wherein  $X^1$  and  $X^2$  together with  $Q^2$  to which they are attached form a ring system of formula Ia, then the compound of formula IIIb represents  $HQ^1(X^3)X^4$  where  $X^3$  and  $X^4$  together with  $Q^1$  to which they are attached form a ring system of formula Ib. Preferably, the ring system of formula Ia is the same as the ring system of formula Ib.


Most preferably, the compound of formula IIIa is identical to the compound of formula IIIb.

20


A compound of formula II, where  $LG_2$  represents hydroxyl or  $NR^{23}R^{24}$ , may be prepared by reaction of a compound of formula IV, wherein  $A_1$ ,  $LG_1$ ,  $K$ ,  $D$ ,  $E$ ,  $M$ ,  $L_1$ ,  $L_2$ ,  $n$  and  $m$  are as defined for a compound of formula II, and  $Li$  represents lithium,

25

54



with a compound of formula Va



(Va)

5

wherein  $A_2$  is as defined for a compound of formula II and  $LG_2$  represents oxygen (thereby forming a hydroxyl derivative following reaction with compound IV) or  $NR^{23}R^{24}$ .

10 Preferably, in a compound of formula Va,  $A_2$  represents methylene and  $LG_2$  represents  $NMe_2$ . For example, a compound of formula Va may represent Eschenmosers salt  $I^-CH_2N^+Me_2$  (see Glidewell C, Journal of Organometallic Chemistry, 527, (1997), p.259-261).

15


Alternatively, in a compound of formula Va,  $A_2$  represents methylene and  $LG_2$  represents oxygen. For example, a compound of formula Va may represent formaldehyde, which for practical purposes may be paraformaldehyde.

20

The reaction may be carried out using methods which are well known to those skilled in the art. For example, the reaction may be accomplished by stirring a solution of the compound of formula IV and Va in an appropriate solvent, 25 such as diethyl ether, at room temperature.

Preferably, in a compound of formula IV,  $LG_1$  does not represent hydroxyl, as the hydroxyl functionality will typically have to be protected prior to ortho-lithiating the precursor compound (compound VI below) to the compound of formula IV. Preferably, in a compound of formula IV,  $LG_1$  represents  $NR^{23}R^{24}$ , most preferably  $LG_1$  represents  $NMe_2$ .

A compound of formula IV may be prepared by ortho-lithiation of a compound of formula VI, wherein  $A_1$ ,  $LG_1$ ,  $K$ ,  $D$ ,  $E$ ,  $M$ ,  $L_1$ ,  $L_2$ ,  $n$  and  $m$  are as defined for a compound of formula IV,




by reaction with an alkyl lithium (e.g. n-butyl lithium). Preferably, in a compound of formula VI,  $LG_1$  does not represent hydroxyl, as the hydroxyl functionality will typically have to be protected prior to performing the ortho-lithiation reaction. Preferably, in a compound of formula VI,  $LG_1$  represents  $NR^{23}R^{24}$ , most preferably  $LG_1$  represents  $NMe_2$ .

Typically, the ortho-lithiation reaction of compounds of formula VI with an alkyl lithium is performed in an inert




solvent, for example tetrahydrofuran or hexane, at low temperatures (e.g.  $-78^{\circ}\text{C}$ ), under a nitrogen atmosphere.

A compound of formula VI, where  $\text{LG}_1$  represents hydroxyl or  $\text{NR}^{23}\text{R}^{24}$ , may be prepared from a compound of formula VII wherein K, D, E, M,  $\text{L}_1$ ,  $\text{L}_2$ , n and m are as defined for a compound of formula VI and Li represents lithium




by reaction with a compound of formula Vb



wherein  $\text{A}_1$  is as defined for a compound of formula VI and  $\text{LG}_1$  represents oxygen (thereby forming a hydroxyl derivative following reaction with compound VII) or  $\text{NR}^{23}\text{R}^{24}$ . Preferably,  $\text{LG}_1$  represents  $\text{NR}^{23}\text{R}^{24}$ , especially  $\text{NMe}_2$ . In other words, a compound of formula Vb is preferably Eschenmosers salt. The reaction may be accomplished using similar conditions as described for the preparation of a compound of formula II above.


Similarly, a compound of formula VII may be prepared by lithiation of a compound of formula VIII




wherein K, D, E, M,  $L_1$ ,  $L_2$ , n and m are as defined for a compound of formula VII.

- 5 Suitably, a compound of formula I wherein K represents  $A_3-Q^3(X^5)X^6$  may be prepared from a compound of formula IX wherein  $A_1$ ,  $A_2$ ,  $A_3$ , D, E, M,  $L_1$ ,  $L_2$ , n and m are as defined for a compound of formula I, and  $LG_1$ ,  $LG_2$  and  $LG_3$  represent suitable leaving groups as defined herein,

10



by reaction with a compound of formula IIIa and IIIb as defined herein, and a compound of formula IIIc



(IIIc)

wherein  $\text{X}^5$  represents  $\text{CR}^{13}(\text{R}^{14})(\text{R}^{15})$ , congressyl or  
 5 adamantyl,  $\text{X}^6$  represents  $\text{CR}^{16}(\text{R}^{17})(\text{R}^{18})$ , congressyl or  
 adamantyl, or  $\text{X}^5$  and  $\text{X}^6$  together with  $\text{Q}^3$  form a 2-phospha-  
 adamantyl group, or  $\text{X}^5$  and  $\text{X}^6$  together with  $\text{Q}^3$  to which  
 they are attached form a ring system of formula Ic; and  
 $\text{R}^{13}$  to  $\text{R}^{18}$ , congressyl, adamantyl,  $\text{Q}^3$ , 2-phospha-adamantyl,  
 10 and the ring system of formula Ic are as defined for a  
 compound of formula I.

Preferably, when the compound of formula IIIa represents  
 $\text{HQ}^2(\text{CR}^1(\text{R}^2)(\text{R}^3))\text{CR}^4(\text{R}^5)(\text{R}^6)$ , then the compound of formula  
 15 IIIb represents  $\text{HQ}^1(\text{CR}^7(\text{R}^8)(\text{R}^9))\text{CR}^{10}(\text{R}^{11})(\text{R}^{12})$  and the  
 compound of formula IIIc represents  
 $\text{HQ}^3(\text{CR}^{13}(\text{R}^{14})(\text{R}^{15}))\text{CR}^{17}(\text{R}^{18})(\text{R}^{19})$ ,  $\text{Q}^1$  is the same as  $\text{Q}^2$  and  $\text{Q}^3$ ,  
 $\text{R}^1$  is the same as  $\text{R}^7$  and  $\text{R}^{13}$ ,  $\text{R}^2$  is the same as  $\text{R}^8$  and  $\text{R}^{14}$ ,  
 $\text{R}^3$  is the same as  $\text{R}^9$  and  $\text{R}^{15}$ ,  $\text{R}^4$  is the same as  $\text{R}^{10}$  and  $\text{R}^{16}$ ,  
 20  $\text{R}^5$  is the same as  $\text{R}^{11}$  and  $\text{R}^{17}$ , and  $\text{R}^6$  is the same as  $\text{R}^{12}$  and  
 $\text{R}^{18}$ .

Preferably, when  $\text{X}^1$  and  $\text{X}^2$  in the compound of formula IIIa  
 independently represents adamantyl, then  $\text{X}^3$  and  $\text{X}^4$  in the  
 25 compound of formula IIIb independently represent  
 adamantyl, and  $\text{X}^5$  and  $\text{X}^6$  in the compound of formula IIIc  
 independently represent adamantyl. Most preferably,  $\text{X}^1$  to  
 $\text{X}^6$  represent the same adamantyl group.

30 Preferably, when  $\text{X}^1$  and  $\text{X}^2$  in the compound of formula IIIa  
 independently represents congressyl, then  $\text{X}^3$  and  $\text{X}^4$  in the  
 compound of formula IIIb independently represent  
 congressyl, and  $\text{X}^5$  and  $\text{X}^6$  in the compound of formula IIIc

independently represent congressyl. Most preferably,  $X^1$  to  $X^6$  represent the same congressyl group.

Preferably, when  $X^1$  and  $X^2$  together with  $Q^2$  to which they  
5 are attached in the compound of formula IIIa form a 2-phospha-adamantyl group, then  $X^3$  and  $X^4$  together with  $Q^1$  in the compound of formula IIIb, and  $X^5$  and  $X^6$  together with  $Q^3$  in the compound of formula IIIc both form a 2-phospha-adamantyl group.

10

Preferably, when  $X^1$  and  $X^2$  together with  $Q^2$  to which they  
are attached form a ring system of formula Ia, then  $X^3$  and  
 $X^4$  together with  $Q^1$  to which they are attached in the  
compound of IIIb forms a ring system of formula Ib, and  $X^5$   
15 and  $X^6$  together with  $Q^3$  to which they are attached in the  
compound IIIc forms a ring system of formula Ic.


Most preferably, the compounds of formula IIIa, IIIb and  
IIIc are identical.

20

Suitably,  $LG_3$  represents a leaving group as defined herein  
in respect of  $LG_1$  and  $LG_2$ . Preferably,  $LG_3$  represents  
 $NR^{23}R^{24}$  or hydroxyl. Most preferably,  $LG_3$  represents  $NMe_2$ ,  
particularly when both  $LG_1$  and  $LG_2$  also represent  $NMe_2$ .

25

Similarly, the compound of formula IX, where  $LG_3$   
represents hydroxyl or  $NR^{23}R^{24}$ , may be prepared by ortho-  
lithiation of a compound of formula II wherein  $A_1$ ,  $A_2$ ,  $LG_1$ ,  
 $LG_2$ ,  $D$ ,  $E$ ,  $M$ ,  $L_1$ ,  $L_2$ ,  $n$  and  $m$  are as defined for a compound  
30 of formula II and  $K$  represents hydrogen, followed by  
reaction with a compound of formula Vc




## (Vc)

wherein  $A_3$  is as defined for a compound of formula IX and  $LG_3$  represents oxygen or  $NR^{23}R^{24}$ . Preferably, when  
 5 employing a compound of formula II to synthesise a compound of formula IX,  $LG_1$  and  $LG_2$  do not represent hydroxyl, as the hydroxyl functionality will typically have to be protected prior to performing the ortho-lithiation reaction. Preferably, both  $LG_1$  and  $LG_2$   
 10 represent  $NR^{23}R^{24}$ , most preferably both  $LG_1$  and  $LG_2$  represent  $NMe_2$ .

Similarly, a compound of formula I wherein K represents  $-A_3-Q^3(X^5)X^6$  and D represents  $-A_4-Q^4(X^7)X^8$  may be prepared  
 15 from a compound of formula IX wherein  $A_1$ ,  $A_2$ ,  $A_3$ ,  $LG_1$ ,  $LG_2$ ,  $LG_3$ , E, M,  $L_1$ ,  $L_2$ , n and m are as defined for a compound of formula IX and D represents hydrogen, by sequential ortho-lithiation and reaction with a compound of formula V(d)


20



## (Vd)

wherein  $A_4$  is as defined for a compound of formula I and  $LG_4$  represents oxygen or  $NR^{23}R^{24}$  as defined herein, to form  
 25 a compound of formula X followed by reaction of the resultant compound of formula X with a phosphine, arsine or stilbene derivative of formula IIIa, IIIb, IIIc as defined herein and a compound of formula IIId

30



## (IIId)

wherein  $X^7$  represents  $CR^{31}(R^{32})R^{33}$ , congressyl or adamantyl,  $X^8$  represents  $CR^{34}(R^{35})(R^{36})$ , congressyl or adamantyl, or  $X^7$  and  $X^8$  together with  $Q^4$  form a 2-phospha-adamantyl group, or  $X^7$  and  $X^8$  together with  $Q^4$  forms a ring system of  
 5 formula Id, and  $R^{31}$  to  $R^{36}$ , adamantyl,  $Q^4$ , 2-phospha-adamantyl, and the ring system of formula Id are as defined for a compound of formula I.

Preferably, when the compound of formula IIIa represents  
 10  $HQ^2(CR^1(R^2)(R^3))CR^4(R^5)(R^6)$ , then the compound of formula IIIb represents  $HQ^1(CR^7(R^8)(R^9))CR^{10}(R^{11})(R^{12})$ , the compound of formula IIIc represents  $HQ^3(CR^{13}(R^{14})(R^{15}))CR^{16}(R^{17})(R^{18})$  and the compound of formula IIId represents  $HQ^4(CR^{31}(R^{32})(R^{33}))CR^{34}(R^{35})(R^{36})$ , and  $Q^1$  is the same as  $Q^2$ ,  $Q^3$   
 15 and  $Q^4$ ,  $R^1$  is the same as  $R^7$ ,  $R^{13}$  and  $R^{31}$ ,  $R^2$  is the same as  $R^8$ ,  $R^{14}$  and  $R^{32}$ ,  $R^3$  is the same as  $R^9$ ,  $R^{15}$  and  $R^{33}$ ,  $R^4$  is the same as  $R^{10}$ ,  $R^{16}$  and  $R^{34}$ ,  $R^5$  is the same as  $R^{11}$ ,  $R^{17}$  and  $R^{35}$ , and  $R^6$  is the same as  $R^{12}$ ,  $R^{18}$  and  $R^{36}$ .

20 Preferably, when  $X^1$  and  $X^2$  in the compound of formula IIIa independently represent adamantyl, then  $X^3$  and  $X^4$  in the compound of formula IIIb independently represent adamantyl,  $X^5$  and  $X^6$  in the compound of formula IIIc independently represent adamantyl, and  $X^7$  and  $X^8$  in the  
 25 compound of formula IIId independently represent adamantyl.

Preferably, when  $X^1$  and  $X^2$  in the compound of formula IIIa independently represent congressyl, then  $X^3$  and  $X^4$  in the  
 30 compound of formula IIIb independently represent congressyl,  $X^5$  and  $X^6$  in the compound of formula IIIc independently represent congressyl, and  $X^7$  and  $X^8$  in the

compound of formula IIIId independently represent congressyl.


Preferably, when  $X^1$  and  $X^2$  together with  $Q^2$  to which they  
5 are attached in the compound of formula IIIa form a 2-phospha-adamantyl group, then  $X^3$  and  $X^4$  together with  $Q^1$  in the compound of formula IIIB,  $X^5$  and  $X^6$  together with  $Q^3$  in the compound of formula IIIC, and  $X^7$  and  $X^8$  together with  $Q^4$  in the compound of formula IIId each independently form  
10 a 2-phospha-adamantyl group.

Preferably, when  $X^1$  and  $X^2$  together with  $Q^2$  to which they  
are attached in the compound of formula IIIa forms a ring  
system of formula Ia, then  $X^3$  and  $X^4$  together with  $Q^1$  in a  
15 compound of formula IIIB,  $X^5$  and  $X^6$  together with  $Q^3$  in a compound of formula IIIC, and  $X^7$  and  $X^8$  together with  $Q^4$  in a compound of formula IIId, form a ring system of formula Ib, Ic and Id, respectively.


20 Most preferably, the compounds of formula IIIa, IIIB, IIIC and IIId are identical.

Suitably, when preparing the 1,2,3,4-substituted derivative of the compound of formula I, preferably  $LG_1$ ,  
25  $LG_2$  and  $LG_3$  of the compound of formula IX do not represent hydroxyl, but each represents  $NR^{23}R^{24}$  as defined herein.

Similarly, a compound of formula I wherein K represents  $-A_3-Q^3(X^5)X^6$ , D represents  $-A_4-Q^4(X^7)X^8$ , and E represents  
30  $-A_5-Q^5(X^9)X^{10}$  may be prepared from a compound of formula XI




wherein  $A_1$ ,  $A_2$ ,  $A_3$ ,  $LG_1$ ,  $LG_2$ ,  $LG_3$ ,  $M$ ,  $L_1$ ,  $L_2$ ,  $n$  and  $m$  are as defined for a compound of formula IX,  $A_4$  is as defined for  
 5 a compound of formula I,  $LG_4$  represents a leaving group, and  $E$  represents hydrogen, by sequential ortho-lithiation and reaction with a compound of formula V(e)



10

wherein  $A_5$  is as defined for a compound of formula I and  $LG_5$  represents oxygen or  $NR^{23}R^{24}$  as defined herein, followed by reaction of the resultant compound with a  
 15 phosphine, arsine or stilbene derivative of formula IIIa, IIIb, IIIc, IIId as defined herein and formula IIIe



20

wherein  $X^9$  represents  $CR^{37}(R^{38})(R^{39})$ , congressyl or adamantyl,  $X^{10}$  represents  $CR^{40}(R^{41})(R^{42})$ , congressyl or adamantyl, or  $X^9$  and  $X^{10}$  together with  $Q^5$  form a 2-phospha-adamantyl group, or  $X^9$  and  $X^{10}$  together with  $Q^5$  form a ring  
 25 system of formula Ie, and  $R^{37}$  to  $R^{42}$ , adamantyl, 2-phospha-



adamantyl,  $Q^5$  and the ring system of formula Ie are as defined for a compound of formula I.

Suitably, when preparing the 1,2,3,4,5-substituted  
5 derivative of formula I, preferably  $LG_1$ ,  $LG_2$ ,  $LG_3$  and  $LG_4$  of the compound of formula X do not represent hydroxyl, but each represents  $NR^{23}R^{24}$  as defined herein.


Preferably, the compounds of formula IIIa, IIIb, IIIc,  
10 IIId and IIIe are identical.

Compounds of formula IIIa wherein  $X^1$  and  $X^2$  together with  $Q^2$  to which they are attached form a 2-phospha-adamantyl group may be prepared by methods well known to those  
15 skilled in the art. Suitably, certain 2-phospha-adamantyl compounds are obtainable from Cytec Canada Inc of 901 Garner Road, Niagara Falls, Ontario, Canada L2E 6T4. Likewise corresponding 2-phospha-adamantyl compounds of formula IIIb, IIIc, IIId and IIIe may be obtained from the  
20 same supplier or prepared by analogous methods.

Compounds of formula IIIa where  $X^1$  and  $X^2$  represent adamantyl may be prepared by methods well known to those skilled in the art, for example by reacting adamantane (or  
25 substituted derivative thereof as defined herein) with phosphorous trichloride and aluminium chloride, followed by reduction of the intermediate  $(adamantyl)_2-P(O)Cl$  derivative. Likewise corresponding compounds of formula IIIb, IIIc, IIId and IIIe where  $X^2$  to  $X^{10}$  respectively  
30 represent adamantyl may be prepared by analogous methods.

Compounds of formula IIIa where  $X^1$  and  $X^2$  together with  $Q^2$  to which they are attached form a ring system of formula


Ia may be prepared by reacting a phosphine ( $\text{PH}_3$ ) with a compound of formula XII




where  $\text{Y}^1$  and  $\text{R}^{49}$  to  $\text{R}^{55}$  are as defined for a compound of formula I. Corresponding compounds of formula IIIb, IIIc, IIId and IIIe may be synthesised by analogous methods.

10

Conveniently, a compound of formula I wherein K, D, E, M,  $\text{A}_2$ ,  $\text{A}_1$ ,  $\text{L}_2$ ,  $\text{L}_1$ ,  $\text{Q}^1$ ,  $\text{Q}^2$ , m and n are as defined for a compound of formula I and  $\text{X}^1$  and  $\text{X}^2$  together with  $\text{Q}^2$  to which they are attached form a ring system of formula Ia and  $\text{X}^3$  and  $\text{X}^4$  together with  $\text{Q}^1$  to which they are attached form a ring system of formula Ib, may be formed by reacting a compound of formula XV



wherein K, D, E, M, A<sub>2</sub>, A<sub>1</sub>, L<sub>2</sub>, L<sub>1</sub>, Q<sup>1</sup>, Q<sup>2</sup>, m and n are as defined for a compound of formula I, with a compound of formula XVIa and XVIb




5

wherein Y<sup>1</sup>, Y<sup>2</sup>, R<sup>49</sup> to R<sup>55</sup> are as defined for a compound of formula I.




Suitably, the reaction may be accomplished by heating the reactants at 120°C for approximately 20 hours.

Thus according to a fifth aspect, the present invention provides a compound of formula XV as defined herein.

Similarly, a compound of formula I wherein D, E, M, A<sub>2</sub>, A<sub>1</sub>, L<sub>2</sub>, L<sub>1</sub>, Q<sup>1</sup>, Q<sup>2</sup>, m and n are as defined for a compound of formula I, X<sup>1</sup> and X<sup>2</sup> together with Q<sup>2</sup> to which they are attached form a ring system of formula Ia, X<sup>3</sup> and X<sup>4</sup> together with Q<sup>1</sup> to which they are attached form a ring system of formula Ib, and K represents -A<sub>3</sub>-Q<sup>3</sup>(X<sup>5</sup>)X<sup>6</sup> wherein X<sup>5</sup> and X<sup>6</sup> together with Q<sup>3</sup> to which it is attached form a ring system of formula Ic and A<sub>3</sub> is as defined for a compound of formula I, may be formed by reacting a compound of formula XVII



wherein D, E, M, A<sub>2</sub>, A<sub>1</sub>, M, L<sub>2</sub>, L<sub>1</sub>, Q<sup>1</sup>, Q<sup>2</sup>, Q<sup>3</sup>, m and n are  
 as defined for a compound of formula I, with a compound of  
 5 formula XVIa, XVIb, XVIc




wherein Y<sup>1</sup>, Y<sup>2</sup>, Y<sup>3</sup>, R<sup>49</sup> to R<sup>55</sup> are as defined for a compound  
 of formula I.

10

Preferably, Y<sup>1</sup> is the same as Y<sup>2</sup> and Y<sup>3</sup> in a compound of  
 formula XVIa, XVIb and XVIc. Most preferably, the  
 compounds of formula XVIa, XVIb and XVIc are identical.

15


Suitably, the compound of formula XV may be prepared from  
 a compound of formula XVIII, where K, D, E, M, L<sub>2</sub>, L<sub>1</sub>, A<sub>1</sub>,  
 A<sub>2</sub>, Q<sup>1</sup>, Q<sup>2</sup>, n and m are as defined for a compound of  
 formula XV,



(XVIII)

by reduction with for example, sodium metabisulfite.

Suitably the compound of formula XVIII may be prepared  
 5 from a compound of formula XIX wherein K, D, E, M, A<sub>2</sub>, A<sub>1</sub>,  
 L<sub>2</sub>, L<sub>1</sub>, m and n are as defined for a compound of formula  
 XVIII



(XIX)

by reaction with an alkyl iodide, such as methyl iodide,  
 10 to form the 1,2-bis-methyl iodide salt derivative of the  
 compound of formula XIX followed by reaction with tris-  
 hydroxymethyl phosphine/stilbene/arsine.

It will be appreciated by those skilled in the art that 1, 2, 3, 4 and 1, 2, 3, 4, 5 substituted compounds may be prepared by analogous methods.

5 The compounds of formula IIIa, IIIb, IIIc, IIId, IIIe, Va, Vb, Vc, Vd, Ve, VIII, XV, XVIa, XVIb, XVIc, XVII, XIX and derivatives thereof, when neither commercially available nor subsequently described, may be obtained using conventional synthetic procedures in accordance with  
10 standard text books on organic chemistry or literature precedent, from readily accessible starting materials using appropriate reagents and reaction conditions.

It will be appreciated by those skilled in the art that,  
15 within certain of the processes described, the order of the synthetic steps employed may be varied and will depend *inter alia* on factors such as the nature of other functional groups present in a particular substrate, the availability of key intermediates and the protecting group  
20 strategy (if any) to be adopted. Clearly, such factors will also influence the choice of reagent for use in the said synthetic steps.

It will also be appreciated that various standard  
25 substituents or functional group interconversions and transformations within certain compounds of formula I will provide other compounds of formula I.

According to a sixth aspect, the present invention  
30 provides a process for the carbonylation of an ethylenically unsaturated compound comprising contacting an ethylenically unsaturated compound with carbon monoxide

and a coreactant in the presence of a compound of the present invention.

Suitable coreactants include compounds comprising a nucleophilic moiety and a mobile hydrogen atom. Thus the compounds of the invention may catalyse hydroformylation, hydrocarboxylation, hydroesterification and hydroamidation reactions of an ethylenically unsaturated compound.

Preferred coreactants include molecular hydrogen, water, alcohols, primary or secondary amines or amides, such as diethylamine, N,N-dimethylethylene diamine, carboxylic acids for example acetic acid, propionic acid and pivalic acid, and aromatic alcohols.

15

Preferably, the coreactant has a hydroxyl functional group or is molecular hydrogen. The hydroxyl containing compounds may be branched or linear, and comprises an alkanol, particularly a C<sub>1</sub>-C<sub>30</sub> alkanol, such as neopentyl alcohol, ethylhexyl alcohol, tert-amyl alcohol, including aryl-alkanols, which may be optionally substituted with one or more substituents selected from lower alkyl, aryl, Het, halo, cyano, nitro, OR<sup>19</sup>, OC(O)R<sup>20</sup>, C(O)R<sup>21</sup>, C(O)OR<sup>22</sup>, NR<sup>23</sup>R<sup>24</sup>, C(O)NR<sup>25</sup>R<sup>26</sup>, C(S)R<sup>27</sup>R<sup>28</sup>, SR<sup>29</sup> or C(O)SR<sup>30</sup> as defined herein. Highly preferred alkanols are C<sub>1</sub>-C<sub>8</sub> alkanols such as methanol, ethanol, propanol, iso-propanol, iso-butanol, t-butyl alcohol, n-butanol, octanol, phenol and chlorocapryl alcohol. Although the monoalkanols are most preferred, polyalkanols, preferably, selected from di-octanols such as diols, triols, tetra-ols and sugars are also possible. Typically, such polyalkanols are selected from 1,2-ethanediol, 1,3-propanediol, glycerol, 1,2,4 butanetriol, 2-(hydroxymethyl)-1,3-propanediol, 1,2,6

trihydroxyhexane, pentaerythritol, 1,1,1  
tri(hydroxymethyl)ethane, nannose, sorbase, galactose and  
other sugars. Preferred sugars include sucrose, fructose  
and glucose. Especially preferred alkanols are methanol  
5 and ethanol.

The amount of alcohol is not critical. Generally, amounts  
are used in excess of the amount of ethylenically  
unsaturated compound to be carbonylated. Thus the alcohol  
10 may serve as the reaction solvent as well, although, if  
desired, separate solvents may also be used.

It will be appreciated that the end product of the  
reaction is determined at least in part by the source of  
15 coreactant compound used. If water is used as the hydroxyl  
group containing compound then the end product is the  
corresponding carboxylic acid, whereas use of an alkanol  
produces the corresponding ester.

20 Preferably, the ethylenically unsaturated compound  
includes from 2 to 20 carbon atoms. More preferably, the  
ethylenically unsaturated compound includes 2 to 14 carbon  
atoms.

25 Suitably, the ethylenically unsaturated compound may  
include more than one carbon-carbon double bond, wherein  
the double bonds are conjugated or non-conjugated.

Preferably, the ethylenically unsaturated compound has 1  
30 to 3 carbon-carbon double bonds per molecule, particularly  
1 to 2 carbon-carbon double bonds, especially only 1  
carbon-carbon double bond per molecule.



Unless otherwise specified, the ethylenically unsaturated compound may, when there are a sufficient number of carbon atoms, be linear or branched, be cyclic, acyclic or part cyclic/acyclic, and/or be substituted or terminated by one  
5 or more substituents selected from lower alkyl, aryl, alkylaryl, Het, alkylHet, halo, OR<sup>19</sup>, OC(O)R<sup>20</sup>, C(O)R<sup>21</sup>, C(O)OR<sup>22</sup>, NR<sup>23</sup>R<sup>24</sup>, C(O)NR<sup>25</sup>R<sup>26</sup>, NO<sub>2</sub>, CN, SR<sup>27</sup> wherein R<sup>19</sup> to R<sup>27</sup> each independently represent hydrogen or lower alkyl. Olefins thus substituted include styrene and alkyl esters  
10 of unsaturated carboxylic acids, such as methacrylate. Suitably, the ethylenically unsaturated compound may exhibit cis (E) and trans (Z) isomerism.

Examples of suitable ethylenically unsaturated compounds  
15 having 2 or more carbon atoms include ethene, propene, but-1-ene, but-2-ene, isobutene, pentenes, hexenes, octenes, dodecenes, 1,5-cyclooctadiene, cyclododecene, methyl pentenoate, pentene nitrile, 1,3 butadiene, 1,3 pentadiene, vinyl acetate and 1,3, hexadiene.  
20 Particularly preferred ethylenically unsaturated compounds include ethene, oct-1-ene, vinyl acetate and 1,3 butadiene, especially ethene.

The process according to the invention may be especially  
25 advantageous for the carbonylation of ethylenically unsaturated compounds which are internally unsaturated, such as but-2-ene, pent-2-ene nitrile, oct-2-ene, oct-3-ene, oct-4-ene or methyl pent-3-enoate. For these compounds side reactions typically occur more readily and  
30 linear products may be more difficult to obtain. Conveniently, the compounds of the invention may permit high regioselectivity towards a linear product following

carbonylation of internally unsaturated ethylenic compounds.

Preferably, the carbonylation process is carried out at a temperature of from 0°C to 250°C, more preferably 40°C to 150°C, most preferably 70°C to 120°C.

Suitably, the carbonylation process is typically carried out at a pressure of at least atmospheric pressure. Preferably, the carbonylation process is performed under a total pressure of greater than or equal to  $1 \times 10^5 \text{Nm}^{-2}$ , more preferably greater than or equal to  $5 \times 10^5 \text{Nm}^{-2}$ , most preferably greater than or equal to  $10 \times 10^5 \text{Nm}^{-2}$ . Preferably, the carbonylation process is performed under a total pressure of less than or equal to  $100 \times 10^5 \text{Nm}^{-2}$ , more preferably less than or equal to  $65 \times 10^5 \text{Nm}^{-2}$ , most preferably less than or equal to  $50 \times 10^5 \text{Nm}^{-2}$ .

Carbon monoxide partial pressures in the range of 1 to 65, particularly 5 to  $50 \times 10^5 \text{Nm}^{-2}$ , are preferred. In the process according to the present invention, the carbon monoxide may be used in pure form or diluted with an inert gas such as nitrogen, carbon dioxide or a noble gas such as argon. Small amounts of hydrogen, typically less than 5% by volume, may also be present.

The ratio (volume/volume) of ethylenically unsaturated compound to coreactant containing compound may vary between wide limits and suitably lies in the range of 1:0.1 to 1:10, preferably from 2:1 to 1:2.

The amount of the catalyst of the invention used in the carbonylation process of the ethylenically unsaturated

compound is not critical. Good results may be obtained by using  $10^{-7}$  to  $10^{-1}$  moles, more preferably  $10^{-6}$  to  $10^{-2}$  moles, of Group VIIIB metal or a compound thereof (referred to as (a) herein) per mole of ethylenically  
5 unsaturated compound.

Preferably, as mentioned herein, the process is performed by the inclusion of a source of anions as defined herein in addition to the compounds of the invention. Suitably, a  
10 source of anions separate from the compounds of the invention may be added to the carbonylation process. Preferably, as mentioned herein, the compounds of the invention include a source of anions. Suitably, the mole ratio of anions to the moles of Group VIIIB metal in the  
15 compounds of the invention when used to carbonylate ethylenically unsaturated compounds lies between wide limits and suitably lies between 2:1 to 2000:1, preferably 10:1 to 200:1.

20 Suitably, the carbonylation of an ethylenically unsaturated compound as defined herein may be performed in one or more aprotic solvents. Suitable solvents include ketones, such as for example methylbutylketone; ethers, such as for example anisole (methyl phenyl ether), 2,5,8-  
25 trioxanonane (diglyme), diethylether, tetrahydrofuran, diphenylether, diisopropylether and the dimethylether of di-ethylene-glycol; esters, such as for example methylacetate, dimethyladipate and butyrolactone; amides, such as for example dimethylacetamide and N-  
30 methylpyrrolidone; and sulfoxides and sulphones, such as for example dimethylsulphoxide, di-isopropylsulphone, sulfolane (tetrahydrothiophene-2,2-dioxide) 2-methylsulfolane and 2-methyl-4-ethylsulfolane.

Very suitable are aprotic solvents having a dielectric constant that is below a value of 50, more preferably in the range of 3 to 8, at 298.15 K and  $1 \times 10^5 \text{ Nm}^{-2}$ . In the present context, the dielectric constant for a given solvent is used in its normal meaning of representing the ratio of the capacity of a condenser with that substance as dielectric to the capacity of the same condenser with a vacuum for dielectric. Values for the dielectric constants of common organic liquids can be found in general reference books, such as the Handbook of Chemistry and Physics, 76<sup>th</sup> edition, edited by David R. Lide et al, and published by CRC press in 1995, and are usually quoted for a temperature of about 20°C or 25°C, i.e. about 293.15K or 298.15 K, and atmospheric pressure, i.e. about  $1 \times 10^5 \text{ Nm}^{-2}$ , or can readily be converted to that temperature and pressure using the conversion factors quoted. If no literature data for a particular compound is available, the dielectric constant may be readily measured using established physico-chemical methods.

For example, the dielectric constant of anisole is 4.3 (at 294.2 K), of diethyl ether is 4.3 (at 293.2 K), of sulfolane is 43.4 (at 303.2 K), of methylpentanoate is 5.0 (at 293.2 K), of diphenylether is 3.7 (at 283.2 K), of dimethyladipate is 6.8 (at 293.2 K), of tetrahydrofuran is 7.5 (at 295.2 K), of methylnonanoate is 3.9 (at 293.2 K). A preferred solvent is anisole.

If the coreactant compound is an alkanol, a further preferred aprotic solvent is the ester carbonylation product of the ethylenically unsaturated compound, carbon monoxide and the alkanol.

The process may advantageously be carried out in an excess of aprotic solvent, i.e. at a ratio (v/v) of aprotic solvent to coreactant containing compound of at least 1:1. Preferably, this ratio ranges from 1:1 to 10:1 and more preferably from 1:1 to 5:1. Most preferably the ratio (v/v) ranges from 1.5:1 to 3:1.

The catalyst compounds of the present invention may act as a "heterogeneous" catalyst or a "homogeneous" catalyst.

By the term "homogeneous" catalyst we mean a catalyst, i.e. a compound of the invention, which is not supported but is simply admixed or formed in-situ with the reactants of the carbonylation reaction (e.g. the ethylenically unsaturated compound, the hydroxyl containing compound and carbon monoxide), preferably in a suitable solvent as described herein.

By the term "heterogeneous" catalyst we mean a catalyst, i.e. the compound of the invention, which is carried on a support.

Thus according to a seventh aspect, the present invention provides a catalyst system comprising a support, preferably an insoluble support, and a compound of the invention as defined herein. Conveniently, the use of an insoluble support permits easy separation of the catalyst, for example by filtration, from the reaction medium.

30

Preferably, the support comprises a polymer such as a polyolefin, polystyrene and polystyrene/divinylbenzene copolymer; a silicon derivative such as a functionalised

silica, a silicone or a silicone rubber; or other porous particulate material such as for example inorganic oxides and inorganic chlorides such as alumina and montmorillonite.

5

Preferably the support material is porous silica which has a surface area in the range of from 10 to 700 m<sup>2</sup>/g, a total pore volume in the range of from 0.1 to 4.0 cc/g and an average particle size in the range of from 10 to 500 μm.

10 More preferably, the surface area is in the range of from 50 to 500 m<sup>2</sup>/g, the pore volume is in the range of from 0.5 to 2.5 cc/g and the average particle size is in the range of from 20 to 200 μm. Most desirably the surface area is in the range of from 100 to 400 m<sup>2</sup>/g, the pore  
15 volume is in the range of from 0.8 to 3.0 cc/g and the average particle size is in the range of from 30 to 100 μm. The average pore size of typical porous support materials is in the range of from 10 to 1000 Å. Preferably, a support material is used that has an average  
20 pore diameter of from 50 to 500 Å, and most desirably from 75 to 350 Å. It may be particularly desirable to dehydrate the silica at a temperature of from 100°C to 800°C anywhere from 3 to 24 hours.

25 Suitably, the support may be flexible or a rigid support, the insoluble support is coated and/or impregnated with the compound of the invention by techniques well known to those skilled in the art. Alternatively, the compound of the invention is fixed to the surface of insoluble  
30 support, optionally via a covalent bond, and optionally includes a bifunctional spacer molecule to space the compound from the insoluble support.

The compounds of the invention may be fixed to the surface of the insoluble support by promoting reaction of a functional group present in the compound of formula I, for example a substituent of the ligand  $L_1$  or a substituent K, D and E of the cyclopentadienyl moiety, with a complimentary reactive group present on or previously inserted into the support. The combination of the reactive group of the support with a complimentary substituent of the compound of the invention provides a heterogeneous catalyst where the compound of the invention and the support are linked via a linkage such as an ether, ester, amide, amine, urea, keto group.

The choice of reaction conditions to link a compound of the present invention to the support depend upon the nature of the substituents(s) of the compound and the groups of the support. For example, reagents such as carbodiimides, 1,1'-carbonyldiimidazole, and processes such as the use of mixed anhydrides, reductive animation may be employed.

According to an eighth aspect, the present invention provides the use of a compound of the invention or a compound of the invention attached to a support as a catalyst.

It will be appreciated that any of the features set forth in the first aspect of the invention may be regarded as preferred features of the second, third, fourth, fifth, sixth, seventh and eighth aspect of the present invention and vice versa.

The invention will now be described by way of the following non-limiting examples.

Example 1

5 Preparation of 1,2-bis-(dimethylaminomethyl)ferrocene

n-Butyllithium (Aldrich, 2.5 molar in hexane, 24 ml, 54 mmol) is added to a solution of (dimethylaminomethyl)ferrocene (Aldrich, 13.13 g, 10.69  
10 ml, 48.97 mmol) in diethyl ether (80 ml) under nitrogen at a temperature of 25°C and the reaction mixture stirred for 4 hours. The resulting red solution is then cooled to approximately -70°C in a dry ice/acetone bath and Eschenmosers salt (ICH<sub>2</sub>NMe<sub>2</sub>) (Aldrich, 10 g, 54 mmol) is  
15 added. The reaction is allowed to warm to room temperature and stirred overnight.

The resultant solution is quenched with excess aqueous sodium hydroxide and the resulting product extracted with  
20 diethyl ether (3 x 80 ml) dried over anhydrous magnesium sulfate, filtered over celite, and volatiles removed in vacuo to yield the crude title compound as a light orange crystalline solid. The crude product is recrystallised from light petrol with cooling to -17°C and the  
25 recrystallised product washed with cold petrol to yield the title compound as a light orange solid (13.2 g, 74%). The compound can be further purified by sublimation to give 8.5 g (52%) of the title compound (mpt 74°C).

30 <sup>1</sup>H NMR(250 MHz; CDCl<sub>3</sub>): δ4.23(brd, 2H); 4.11-4.10(t, 1H); 4.04(s, 5H); 3.43, 3.38, 3.23, 3.18 (AB quartet, 2H); 2.22(s, 6H).



$^{13}\text{C}$  NMR (63 MHz;  $\text{CDCl}_3$ ):  $\delta$ 83.81; 70.40; 69.25; 66.84; 57.35; 45.23.

Elemental analysis: Found: C 63.7%; H 8.9%; N 9.5%  
5 Calculated: C 64.0%; H 8.1%; N 9.4%

### Example 2

#### Preparation of 1,2-bis-(ditertbutylphosphinomethyl)ferrocene

10 Di-tertbutylphosphine (Aldrich, 0.616 ml, 3.33 mmol) was added to a solution of 1,2-bis(dimethylaminomethyl)ferrocene (Example 1, 0.5 g, 1.66 mmol) in anhydrous acetic acid (100 ml) under nitrogen and the resulting mixture is stirred at 80°C for 72 hours. The  
15 anhydrous acetic acid is removed *in vacuo* at approximately 70°C to yield the crude title product as an orange/yellow solid. The crude product is recrystallised from ethanol with cooling to -17°C, filtered and the filtrate washed with cold ethanol to yield the title compound as a pale  
20 yellow solid (0.365 g, 44%, 84°C).

$^1\text{H}$  NMR (250 MHz;  $\text{CDCl}_3$ ):  $\delta$ 4.4 (2H, d,  $J = 2\text{Hz}$ ); 3.95 (5H, s); 3.75 (1H, t, 2Hz); 2.8 (2H, dd, 12Hz, 2Hz); 2.6 (2H, dd, 12Hz, 2Hz); 1.1 (36H, m).

25

$^{13}\text{C}$  NMR (63 MHz;  $\text{CDCl}_3$ ):  $\delta$ 86.73 (d, 5.46 Hz); 70.08 (d, 4.41 Hz); 69.4665(s); 63.75(s); 31.80 (d, 2Hz); 31.45 (d, 1.98Hz); 29.89 (d, 1.88 Hz).

30  $^{31}\text{P}$  NMR (101 MHz;  $\text{CDCl}_3$ ):  $\delta$ 15.00 ppm.

Elemental analysis: Found: C:66.79%; H:9.57%  
Calculated: C:66.93%; H:9.63%

Example 3Preparation of 1-hydroxymethyl-2-dimethylaminomethyl ferrocene

5

n-Butyl lithium (Aldrich, 1.6 molar in diethyl ether, 5.14 ml, 8.24 mmol) is added to a solution of 1-dimethylaminomethyl ferrocene (Aldrich, 1.0g, 4.12mmol) in diethyl ether (20mL) under argon. The reaction is stirred for 3 hours and develops a reddish colour. The solution is then cooled in a dry ice/acetone bath, calcined para-formaldehyde (0.247g, 2 times excess) added and the resultant mixture stirred overnight at room temperature. The reaction is then quenched with water, extracted with diethyl ether, dried over  $\text{MgSO}_4$ , and filtered over celite. The solvent is removed in vacuo to yield crude title compound. The crude product is applied to a neutral alumina column, which is eluted with petrol/diethyl ether (9:1 ratio) to remove the starting material, 1-dimethylaminomethyl ferrocene. The column is then eluted with substantially pure ethyl acetate to elute the title compound. The ethyl acetate is removed in vacuo, to yield the title compound as an orange oil/crystalline mass.

25  $^1\text{H}$  NMR (250 MHz;  $\text{CDCl}_3$ )  $\delta$ 2.131 (s, 6 H),  $\delta$ 2.735 (d, 1 H, 12.512 Hz),  $\delta$ 3.853 (d, 1 H, 12.512 Hz),  $\delta$ 3.984 (dd, 1 H, 2.156 Hz),  $\delta$ 4.035 (s, 5 H),  $\delta$ 4.060 (dd, 1 H, 2.136 Hz)  $\delta$ 4.071 (d, 1 H, 12.207 Hz),  $\delta$ 4.154 (m, 1 H),  $\delta$ 4.73 (d, 1 H, 12.207 Hz).

30

$^{13}\text{C}$  NMR (61 MHz;  $\text{CDCl}_3$ )  $\delta$ 7.688,  $\delta$ 84.519,  $\delta$ 70.615,  $\delta$ 68.871,  $\delta$ 68.447,  $\delta$ 65.369,  $\delta$ 60.077,  $\delta$ 58.318,  $\delta$ 44.414

COSY 2D  $^1\text{H}$  NMR

Partly obscured doublet at 4.071ppm and its coupling to the doublet at 4.73 ppm confirmed.

5 Infrared spectra ( $\text{CHCl}_3$ ) (c.a. 0.06g / 0.8mL)

2953.8  $\text{cm}^{-1}$  , 2860.6  $\text{cm}^{-1}$ , 2826.0  $\text{cm}^{-1}$ , 2783.4  $\text{cm}^{-1}$ , 1104.9  $\text{cm}^{-1}$

Example 4

10 Preparation of 1,2-bis-(ditertbutylphosphinomethyl)ferrocene

Di-tertbutylphosphine (Aldrich, 0.54 ml, 2.93 mmol) is added to a solution of 1-hydroxymethyl-2-dimethylaminomethyl ferrocene (Example 3, 0.2 g, 0.753 mmol) in anhydrous acetic acid (15 ml) and acetic anhydride (0.753 mmol) under argon and the resulting mixture is stirred at 80°C for 72 hours. The anhydrous acetic acid is removed *in vacuo* at approximately 70°C to yield the crude title product as an orange/yellow solid.

15 20 The crude product is recrystallised from ethanol with cooling to -17°C, filtered and the filtrate washed with cold ethanol to yield the title compound as an orange solid (0.23 g)

25  $^1\text{H}$  NMR (250 MHz;  $\text{CDCl}_3$ )  $\delta$ 4.351 (d, 2 H, 2Hz),  $\delta$ 4.022 (s, 5 H),  $\delta$ 3.827 (t, 1 H, 2 Hz),  $\delta$ 2.858 (ddd, 2 H,  $J_{\text{HH}}$  15.869 Hz,  $J_{\text{HP1}}$  3.320 Hz,  $J_{\text{HP2}}$  1.831 Hz),  $\delta$ 2.679 (dd, 2 H,  $J_{\text{HH}}$  15.869 Hz,  $J_{\text{HP}}$  2.441 Hz),  $\delta$ 1.166 (d, 18 H, 12.817 Hz),  $\delta$ 1.123 (d, 18 H, 12.512 Hz)

30

FTIR (Chloroform, NaCl plates)

1104.1  $\text{cm}^{-1}$ , 2863  $\text{cm}^{-1}$ , 2896.0  $\text{cm}^{-1}$ , 2940.0  $\text{cm}^{-1}$ , 2951.8  $\text{cm}^{-1}$

<sup>31</sup>P NMR (101 MHz; CDCl<sub>3</sub>): δ15.00 ppm.

Elemental analysis: Found: C:66.5%; H:9.6%  
5 Calculated: C:66.9%; H:9.6%

#### Example 5

#### Preparation of 1-hydroxymethyl-2,3-bis- (dimethylaminomethyl)ferrocene

10

To a stirred solution of 1,2-bis-(dimethylaminomethyl)ferrocene (Example 1, 0.70g, 2.32 mmol) in diethyl ether (15 cm<sup>3</sup>) under argon is added 1.2 equivalents n-butyl lithium (Aldrich, 1.75mL, 1.6M in diethyl ether) and the  
15 mixture stirred for three hours to yield a red solution. The reaction mixture is cooled in a dry ice/acetone bath, calcined paraformaldehyde added in 2:1 excess, and the resultant mixture stirred at room temperature overnight. The mixture is quenched with water and extracted with  
20 diethyl ether. The ethereal extracts are dried over MgSO<sub>4</sub>, filtered over celite and the solvent removed in vacuo, to yield the title compound (0.7g , 2.12 mmol, 91%) as an orange oil., which partially crystallized on cooling.

25 <sup>1</sup>H NMR (250 MHz; CDCl<sub>3</sub>) δ 2.133 (s, 6 H ), δ 2.171 (s, 6 H), δ 2.910 (d, 1 H, 12.817 Hz), δ 2.998 (d, 1 H, 12.512 Hz), δ 3.425 (d, 1 H, 12.817 Hz), δ 3.812 (d, 1 H, 12.512 Hz), δ 3.962 (s, 5 H), δ 3.99 (d, 1 H, 12.207 Hz) (partly obscured by large cp-ring peak at δ 3.962), δ 4.068 (d, 1  
30 H, δ2.136 Hz), δ 4.125 )d, 1 H, δ 2.136 Hz), δ 4.747 (d, 1 H, 12.207 Hz)

$^{13}\text{C}$  NMR (60 MHz;  $\text{CDCl}_3$ )  $\delta$  44.529,  $\delta$  45.244,  $\delta$  55.798,  $\delta$  57.906,  $\delta$  60.271,  $\delta$  67.944,  $\delta$  68.277,  $\delta$  69.612,  $\delta$  84.850,  $\delta$  88.322

Infrared spectra ( $\text{CDCl}_3$  / thin film NaCl plates)

5 3380.6  $\text{cm}^{-1}$  (br), 2955.7  $\text{cm}^{-1}$  (m), 2862.6  $\text{cm}^{-1}$ , 2825.9  $\text{cm}^{-1}$  (m), 2774.3  $\text{cm}^{-1}$  (m), 1353.5  $\text{cm}^{-1}$  (m), 1104.9  $\text{cm}^{-1}$  (m), 1038.9  $\text{cm}^{-1}$  (m), 1006.8  $\text{cm}^{-1}$  (s)

Elemental analysis: Found: C: 62.3%; H: 7.8%; N: 8.8%

10 Calculated: C: 61.8%; H: 7.9%; N: 8.5%

#### Example 6

##### Preparation of

##### 1,2,3-tris-(ditertbutylphosphinomethyl)ferrocene

15

Di-tert-butylphosphine (Aldrich, 2.60 mL, 13.98 mmol) and acetic anhydride (0.24 mL, 2.33 mmol) is added to a solution of 1-hydroxymethyl-2,3-bis-(dimethylaminomethyl)ferrocene (Example 5, 0.70g, 2.12  
20 mmol) in acetic acid (freshly distilled from acetic anhydride 25  $\text{cm}^3$ ), under argon. The solution is then stirred at 80°C for 7 days, during which time the solution becomes a dark orange colour. The solvent is then removed *in vacuo* and recrystallisation effected from refluxing  
25 ethanol together with cooling to -17°C overnight to yield the title compound (0.43 g, 0.7 mmol, 31%) as a yellow/orange powder.

$^1\text{H}$  NMR (250 MHz,  $\text{CDCl}_3$ )  $\delta$  1.12 (dd - pseudo triplet, 36 H, 30 12.1 Hz),  $\delta$  1.26 (d, 18H, 10.7 Hz),  $\delta$  2.68 (d, 2 H, 17.7 Hz),  $\delta$  2.95 (s, 2 H),  $\delta$  3.07, (m, 2 H),  $\delta$  4.01 (s, 5 H)  $\delta$  4.33 (s, 2 H)

Infrared spectra ( $\text{CHCl}_3$  / thin film NaCl plates )  
1365.5  $\text{cm}^{-1}$ , 1470.3  $\text{cm}^{-1}$  , 2357.1  $\text{cm}^{-1}$  , 2862.8  $\text{cm}^{-1}$  , 2896.7  
 $\text{cm}^{-1}$  , 2939.1  $\text{cm}^{-1}$

5

#### Example 7

Preparation of 1,2-bis-(dicyclohexylphosphinomethyl)  
ferrocene

10

The title compound was prepared in accordance with the  
procedure of Example 2 employing dicyclohexylphosphine  
(Strem of 48 High Street Orwell, Royston, United Kingdom  
SG8 5QW, 659 mg, 3.33 mmol), 1,2-  
15 bis(dimethylaminomethyl)ferrocene (0.5 g, 1.66 mmol) and  
anhydrous acetic acid (100 ml). Yield 0.421 g.

#### Example 8

Preparation of 1,2-bis-(di-iso-butylphosphinomethyl)  
ferrocene

20

The title compound was prepared in accordance with the  
procedure of Example 2 employing di-iso-butylphosphine  
(Strem 486 mg, 3.33 mmol), 1,2-  
25 bis(dimethylaminomethyl)ferrocene (0.5 g, 1.66 mmol) and  
anhydrous acetic acid (100 ml). Yield 0.372 g.

#### Example 9

Preparation of 1,2-bis-(dicyclopentylphosphinomethyl)  
ferrocene

30

The title compound was prepared in accordance with the  
procedure of Example 2 employing dicyclopentylphosphine

(Strem 566 mg, 3.33 mmol), 1,2-bis(dimethylaminomethyl)ferrocene (0.5 g, 1.66 mmol) and anhydrous acetic acid (100 ml). Yield 0.432 g.

5 Example 10

Preparation of 1,2-bis-(diethylphosphinomethyl) ferrocene

The title compound was prepared in accordance with the procedure of Example 2 employing diethylphosphine (Strem  
10 299 mg, 3.33 mmol), 1,2-bis(dimethylaminomethyl)ferrocene (0.5 g, 1.66 mmol) and anhydrous acetic acid (100 ml). Yield 0.254 g.

Example 11Preparation of 1,2-bis(di-isopropylphosphinomethyl) ferrocene

5 The title compound was prepared in accordance with the procedure of Example 2 employing di-iso-propylphosphine (Digital Speciality Chemicals 392 mg, 3.33 mmol), 1,2-bis(dimethylaminomethyl)ferrocene (0.5 g, 1.66 mmol) and anhydrous acetic acid (100 ml). Yield 0.262 g.

10

Example 12Preparation of 1,2-bis-(dimethylphosphinomethyl) ferrocene

15 The title compound was prepared in accordance with the procedure of Example 2 employing dimethylphosphine (Digital Speciality Chemicals, 206 mg, 3.33 mmol), 1,2-bis(dimethylaminomethyl)ferrocene (0.5 g, 1.66 mmol) and anhydrous acetic acid (100 ml). Yield 0.285 g.

20 Example 13Preparation of 1,2-bis-(diadamantylphosphinomethyl)ferrocene bis-methanesulphonate

25 Di-adamantylphosphine (prepared according to J.R.Goerlich, R.Schmutzler; Phosphorus Sulphur and Silicon; 1995, 102, 211-215, 20.0g, 0.066 mol) was added to a solution of 1,2-bis(dimethylaminomethyl)ferrocene (Example 1, 10 g, 0.033 mol) in anhydrous acetic acid (100 ml) under nitrogen and the resulting mixture is stirred at 80°C for 72 hours. The  
30 orange yellow precipitate which forms is filtered and dried in vacuo at approximately 70°C to yield the title compound as an orange/yellow solid. The title compound is insoluble in a range of organic solvents and it is



therefore purified by conversion to the bis-methanesulphonate salt by addition of excess methanesulphonic acid to a methanol slurry of the crude product. This resulted in complete dissolution of the product salt which was then isolated by removal of the methanol in vacuo followed by washing with ether and drying to give the title compound as a pale yellow solid (14.0g, 54%).

<sup>1</sup>H NMR (250 MHz; CD<sub>3</sub>CN): δ4.57 (2H, d, J = 2Hz); 4.35 (5H, s); 4.27 (1H, t, 2Hz); 3.34 (4H, br); 2.6 (6H, br,); 2.35-2.18 (18H br); 2.16-2.0 (18H, br); 1.92-1.72 (24H, br).

<sup>31</sup>P NMR (101 MHz; CD<sub>3</sub>CN): δ26.58 ppm.

Elemental analysis: Found: C:64.15%; H:7.88% Calculated: C:64.29%; H:7.94%

#### Example 14

#### Preparation of 1,2 bis(di-1-adamantylphosphinomethyl) ferrocene-bis-methane sulphonate

The preparation of this ligand was carried out as follows:

#### 14.1 Preparation of (1-Ad)<sub>2</sub>P(O)Cl

Phosphorous trichloride (83 cm<sup>3</sup>, 0.98 mol) was added rapidly via cannula to a combination of aluminium chloride (25.0 g, 0.19 mol) and adamantane (27.2 g, 0.20 mol) affording a tan suspension. The reaction was heated to reflux. After 10 minutes, a yellow-orange suspension was formed. The reaction was refluxed for a total of 6 hours. The excess PCl<sub>3</sub> was removed by distillation at atmospheric

pressure (BP 75°C). On cooling to ambient temperature, an orange solid was formed. Chloroform (250 cm<sup>3</sup>) was added yielding an orange suspension, which was cooled to 0°C. Water (150 cm<sup>3</sup>) was added slowly: initially the suspension  
5 viscosity increased, but on full addition of water the viscosity lessened. From this point the reaction was no longer kept under an atmosphere of Ar. The suspension was Buchner filtered to remove the yellow-orange solid impurity. The filtrate consisted of a two phase system.  
10 The lower phase was separated using a separating funnel, dried over MgSO<sub>4</sub> and Buchner filtered. The volatiles were removed via rotary evaporation, drying finally *in-vacuo*, affording an off-white powder. Yield 35.0 g, 99%. <sup>31</sup>P NMR: = 85 ppm, 99% pure. FW = 352.85.

15

#### 14.2 Preparation of (1-Ad)<sub>2</sub>PH

LiAlH<sub>4</sub> (2.54 g, 67.0 mmol) was added over 90 minutes to a chilled (-10°C) solution of (1-Ad<sub>2</sub>P(O)Cl (10.00 g, 28.3  
20 mmol) in THF (120 cm<sup>3</sup>). The reaction was allowed to warm to ambient temperature then stirred for 20 hours. The grey suspension was cooled to -10°C. HCl (aq. 5 cm<sup>3</sup> c. HCl in 50 cm<sup>3</sup> degassed water) was added slowly via syringe (initially very slowly due to exotherm of reaction),  
25 yielding a two phase system, with some solid material in the lower phase. Further HCl (~ 5 cm<sup>3</sup> c. HCl) was added to improve the separation of the layers. The upper phase was removed via flat ended cannula, dried over MgSO<sub>4</sub> and filtered via cannula. The volatiles were removed *in-vacuo*  
30 affording the product as a white powder, isolated in the glovebox. Yield 6.00 g, 70%. <sup>31</sup>P NMR: = 17 ppm, 100% pure. FW = 302.44.

14.3 Preparation of 1,2-bis(di-1-adamantylphosphinomethyl) ferrocene-bis-methanesulphonate

The title compound was prepared in accordance with the  
5 procedure exemplified in Example 13.

Example 15

Preparation of 1,2-bis(di-1-(3,5-dimethyladamantyl)phosphinomethyl) ferrocene-bis-methanesulphonate

10

15.1 Di-1-(3,5-dimethyladamantyl) phosphinic chloride  
was prepared in accordance with the method of Example 14.1  
except using 1,3 dimethyladamantane 21.7 g (0.132 mol)  
instead of adamantane, and AlCl<sub>3</sub> (18.5 g, 0.14 mol). Yield  
15 23.5 g FW: 409.08. <sup>31</sup>P NMR: δ: 87 ppm (s).

15.2 Di-1-(3,5-dimethyladamantyl) phosphine  
was prepared as Example 14.2 above except using 25.0 g Di-  
1-(3,5-dimethyladamantyl) phosphinic chloride instead of  
20 di-1-adamantyl phosphinic chloride. Yield 15.7 g, FW:  
358.58.. <sup>31</sup>P NMR: δ: 15.7 ppm (s).

15.3 1,2-bis-(di-1-(3,5-dimethyladamantylphosphinomethyl) ferrocene-bis-methanesulphonate

25

The title compound was prepared in accordance with the  
procedure exemplified in Example 13 except using di-1-  
2 (3,5-dimethyl-adamantyl)phosphine (23.69 g, 0.066 mol)  
instead of di-adamantylphosphine. Yield 15 g.

30

Example 16

Preparation of 1,2-bis(di-1-(4-tert-butyl-adamantyl)phosphinomethyl) ferrocene-bis-methanesulphonate

16.1 Di-1-(4-tert-butyladamantyl) phosphinic chloride  
was prepared as per Di-1-adamantyl phosphinic chloride of  
Example 14.1 except using 4-tert-butyladamantane 25.37 g  
5 (0.132 mol) instead of adamantane, and AlCl<sub>3</sub> (18.5 g, 0.14  
mol). Yield 22.6 g FW: 464.98. <sup>31</sup>P NMR: δ: 87 ppm (s).

16.2 Di-1-(4-tert-butyladamantyl) phosphine  
was prepared as per Di-1-adamantyl phosphine of Example  
10 14.2 except using 13.5 g Di-1-(4-tert-butyladamantyl)  
phosphinic chloride instead of di-1-adamantyl phosphinic  
chloride. Yield 9.4 g, FW: 414.48. <sup>31</sup>P NMR: δ: 18.62 ppm  
(s).

15 16.3 1,2-bis(di-1-(4-tert-butyl-adamantyl)  
phosphinomethyl) ferrocene-bis-methanesulphonate

The title compound was prepared in accordance with the  
procedure exemplified in Example 13 except using di-1-(4-  
20 tert-butyladamantyl)phosphine (27.39 g, 0.066 mol) instead  
of di-adamantyl phosphine. Yield 14.52 g.

#### Example 17

25 Preparation of 1,2-bis-(1-adamantyl tert-butyl-  
phosphinomethyl)ferrocene-bis-methanesulphonate

#### 17.1 1-adamantylphosphonic acid dichloride

This compound was synthesised according to the method of  
30 Olah et al (J. Org. Chem. 1990, 55, 1224-1227).

### 17.2 1-adamantyl phosphine

LiAlH<sub>4</sub> (3.5 g, 74 mmol) was added over 2 hours to a cooled solution (0°C) of 1-adamantylphosphonic acid dichloride (15 g, 59 mmol) in THF (250 cm<sup>3</sup>). The reaction was then allowed to warm to ambient temperature and was stirred for 20 hours. The grey suspension was then cooled (0°C) and HCl (75 cm<sup>3</sup>, 1M) was slowly added via syringe, to afford a two phase system with some solid present in the lower phase. Concentrated HCl (8 cm<sup>3</sup>, 11M) was then added to improve the separation of the two layers. The (upper) THF phase was removed via cannula and dried over magnesium sulphate. After filtration via cannula, the volatiles were removed *in-vacuo* to afford the product.

15

### 17.3 1-adamantyl tert-butyl phosphine

nBuLi (20 cm<sup>3</sup>, 32 mmol 1.6M soln) was added over 1 hour to a cooled solution of 1-adamantyl phosphine (5.0 g, 30 mmol) in THF (100 cm<sup>3</sup>). The solution was allowed to warm to room temperature and stirred for a further 2 hours. The solution was recooled to 0°C and tert-butylchloride (2.78 g, 30 mmol) was added and stirring continued for a further 16 hours at room temperature. The reaction mixture was quenched with water and the aqueous phase extracted with dichloromethane (2 x 50 ml). The organic phase was dried over sodium sulphate and evaporated *in-vacuo* to yield the title compound.

30

17.4 1,2-bis-(-1-adamantyl tert-butyl-phosphinomethyl)  
ferrocene-bis-methanesulphonate

The title compound was prepared in accordance with the  
5 procedure exemplified in Example 13 except using 1-  
adamantyl tert-butyl phosphine (14.78 g, 0.066 mol)  
instead of di-adamantyl phosphine. Yield 9.80 g.

Example 18

10 Preparation of 1,2-bis-(di-1-diamantylphosphinomethyl)  
ferrocene-bis-methanesulphonate

18.1 Diamantane

15 This was synthesised according to the method of Tamara et  
al. Organic Syntheses, CV 6, 378.

18.2 Di-1-(diamantane) phosphinic chloride

20 Prepared as per Di-1-adamantyl phosphinic chloride of  
Example 14.1 except using diamantane 20.0 g (0.106 mol)  
and AlCl<sub>3</sub> (16.0 g, 0.12 mol). Yield 25.5 g FW: 456.5.. <sup>31</sup>P  
NMR: δ: 87 ppm (s).

25 18.3 Di-1-(diamantane) phosphine

Prepared as per Di-1-adamantyl phosphine of Example 14.2  
except using 25.0 g Di-1-(diamantane) phosphinic chloride.  
Yield 14.0 g FW: 406.. <sup>31</sup>P NMR: δ: 16.5 ppm (s).

18.4 1,2-bis-(di-1-diamantylphosphinomethyl)ferrocene-  
bis-methanesulphonate

The title compound was prepared in accordance with the  
5 procedure exemplified in Example 13 except using di-1-  
diamantane phosphine (26.79 g, 0.066 mol) instead of di-  
adamantyl phosphine. Yield 12.5 g.

Example 19

10 Preparation of 1,2-bis-(di-(1,3,5,7-tetramethyl-6,9,10-  
trioxa-2-phospha-adamantylmethyl))ferrocene

1,3,5,7-tetramethyl-2,4,8-trioxa-6-phospha-adamantane  
(obtained from Cytec, 14.0g, 0.066 mol) was added to a  
15 solution of 1,2-bis(dimethylaminomethyl)ferrocene (Example  
1, 10 g, 0.033 mol) in anhydrous acetic acid (100 ml)  
under nitrogen and the resulting mixture is stirred at  
80°C for 72 hours. The anhydrous acetic acid is removed in  
vacuo at approximately 70°C to yield the crude title  
20 product as an orange/yellow solid. This is washed with  
hot methanol to give the product as a mixture of isomers  
as an orange solid. (12.0g, 58%).

<sup>1</sup>H NMR (250 MHz; CDCl<sub>3</sub>): δ 4.25-3.95 (8H, br, m); 3.46 (4H,  
25 br); 1.57-2.0 (8H, br, m); 1.43-1.23 (24H, br m).

<sup>31</sup>P NMR (101 MHz; CDCl<sub>3</sub>): δ -27.41 (br), -29.01 (s), -33.9  
(br) ppm.

30 Elemental analysis: Found: C:57.80%; H:7.35% Calculated:  
C:57.87%; H:7.40%

Example 20Preparation of 1,2-bis-(dimethylaminomethyl)ferrocene-bis methyl iodide

5 Methyl iodide (23.28g, 0.164 mol) is added to a solution of 1,2-bis-(dimethylaminomethyl)ferrocene (Example 1, 20g, 0.082 mol) in degassed methanol (100 ml), and the mixture stirred at room temperature under a nitrogen atmosphere for 24 hours. The resulting precipitate is  
10 removed by filtration, washed with ether and dried to yield the title compound (43.0g).

Elemental analysis: Found: C:36.8%; H:5.1%; N,4.8%

Calculated: C:37.0%; H:5.2%; N,4.8%

15

<sup>13</sup>C NMR (D<sub>2</sub>O): δ53.27, δ53.21, δ53.15, δ64.68, δ71.77, δ73.24, δ74.13, δ74.95

Example 21

20 Preparation of 1,2-bis(dihydroxymethylphosphinomethyl)ferrocene

Potassium hydroxide (8.52g, 0.152 mol) is added to a solution of tetrakis(hydroxymethyl) phosphonium chloride  
25 (Aldrich, 38.54g of 80% w/w aqueous solution, 0.162 mol) in degassed methanol (40 ml), and stirred at room temperature under a nitrogen atmosphere for 1 hour. The resultant mixture is added dropwise to a degassed solution of 1,2-bis-(dimethylaminomethyl)ferrocene-bis-methyl  
30 iodide (Example 20, 19.98g, 52.2 mmol) in methanol (40 ml) under nitrogen at room temperature with stirring. The resultant mixture is refluxed under nitrogen for 20 hours, and the solvent removed in vacuo to form a red



precipitate. Water (30 ml), diethyl ether (85 ml) and triethylamine (35 ml) is added to the precipitate and the solution stirred at room temperature for 1 hour. The aqueous layer is removed and re-extracted with diethyl ether (2 x 30 ml). The combined ethereal extracts are washed with water (3 x 20 ml) dried over sodium sulphate and filtered. The ether is removed in vacuo to yield the crude title compound (14.33g, 94% yield) as a microcrystalline orange solid. The crude product is recrystallised from a warm dichloromethane/methanol solution with the addition of light petroleum and cooling to yield the title compound (10.69g, 70% yield) as yellow-orange crystals.

Elemental analysis: Found: C:48.44%; H:4.12%; N,0.0%  
Calculated: C:48.24%; H:4.02%;  
N,0.0%

$^1\text{H}$  NMR:  $\delta$ 1.75 (s, br),  $\delta$ 2.70 (dd, 2 H,  $J_{\text{HH}}^2$  14.2 Hz,  $J_{\text{HP}}^2$  6.6 Hz),  $\delta$ 2.85 (dd, 2 H,  $J_{\text{HH}}^2$  14.2 Hz,  $J_{\text{HP}}^2$  7.9 Hz),  $\delta$ 3.71 (t, 1 H,  $J_{\text{HH}}$  2.44 Hz),  $\delta$ 3.58 (s, 5 H),  $\delta$ 3.98 (d, 2 H,  $J_{\text{HH}}$  2.40 Hz), 4.06 (m, 8 H).

$^1\text{H}\{^{31}\text{P}\}$  NMR:  $\delta$ 1.75 (s, br),  $\delta$ 2.70 (d, 14.3 Hz),  $\delta$ 2.85 (d, 14.3 Hz),  $\delta$ 4.04 (m, 1 H),  $\delta$ 4.06 (s, 8 H),  $\delta$ 4.08 (s, 5H),  $\delta$ 4.1 (m, 2 H)

$^{13}\text{C}$  NMR:  $\delta$ 23.7 (d,  $J_{\text{PC}}^1$  15.6 Hz),  $\delta$ 63.0 (d,  $J_{\text{PC}}^1$  15.6 Hz),  $\delta$ 66.0 (s),  $\delta$ 67.2 (d,  $J_{\text{PC}}^3$  9.2 Hz),  $\delta$ 69.6 (s),  $\delta$ 82.6 (d,  $J_{\text{PC}}^2$  14.7 Hz)

$^{31}\text{P}$  NMR:  $\delta$ -14.7

Infrared spectra ( $\text{CHCl}_3$  / thin film NaCl plates )

3337.8  $\text{cm}^{-1}$  (st, br), further peaks 1104  $\text{cm}^{-1}$  2929.0  $\text{cm}^{-1}$ ,  
3603.7  $\text{cm}^{-1}$  , 3683.7  $\text{cm}^{-1}$ .

5 Example 22

Preparation of 1,2-bis(diphosphinomethyl)ferrocene

1,2-bis(dihydroxymethylphosphinomethyl)ferrocene (Example  
21, 5.45g, 13.70 mmol) and sodium metabisulfite (5.21g,  
10 27.4 mmol) is added to a two-phase solvent system  
consisting of distilled water (60 ml) and light petroleum  
(60 ml). The mixture is refluxed for 3 hours in air. The  
resultant mixture is cooled stirred and the aqueous layer  
removed. The organic layer is washed with distilled water  
15 and the organic solvent removed in vacuo to yield the  
title compound (2.66g, 70% yield) as an orange crystalline  
solid.

Elemental analysis: Found: C:51.65%; H:5.75%

20 Calculated: C:51.80%, H:5.76%

$^1\text{H}$  NMR (250 MHz;  $\text{CDCl}_3$ ):  $\delta$  2.7-2.8 (m, 4H),  $\delta$  3.17 (m,  
2H),  $\delta$  3.18 (m, 2H),  $\delta$  4.04 (t, 1H,  $J=2.54$  Hz),  $\delta$  4.09 (d,  
5H,  $J_{\text{HP}}$  0.4 Hz),  $\delta$  4.13 (d, 2H,  $J=2.54$  Hz)

25

$^{31}\text{P}$  NMR (101 MHz;  $\text{CDCl}_3$ ):  $\delta$  130.0 (t,  $J_{\text{HP}}$  193.0 Hz)

$^{13}\text{C}$  NMR (60 MHz;  $\text{CDCl}_3$ ):  $\delta$  12.9,  $\delta$  65.6,  $\delta$  67.3,  $\delta$  69.4,  $\delta$   
86.9

30

$^{13}\text{C}$  DEPT NMR ( $\text{CDCl}_3$ ):  $\delta$  12.9 ( $\text{CH}^2$ ),  $\delta$  65.6 (CH),  $\delta$  67.3  
(CH),  $\delta$  69.40 (5 x CH)

FTIR (Chloroform, NaCl plates): 2298.5  $\text{cm}^{-1}$  (strong)

Mass spectrum: Found m/z: 278.0088; Calculated m/z 278.0077

5

Example 23

Preparation of 1,2-bis- $\alpha,\alpha$ -(P-(2,2,6,6,-tetramethylphosphinan-4-one))dimethylferrocene

10 2,6-Dimethyl-2,5-heptadiene-4-one (14.6g, 0.106 mol) is added to 1,2-bis-(diphosphinomethyl)ferrocene (Example 22, 14.7g, 0.053 mol) and the mixture heated to 120°C under nitrogen for 20 hours. The reaction mixture is cooled, the crude title compound removed by filtration, washed with  
15 pentene (20 ml) and dried in vacuo to yield the title compound as a yellow-orange solid (24.9g, 85% yield). The title compound was characterised by  $^{31}\text{P}$  NMR and mass spectrum.

20  $^1\text{H}$  NMR (250 MHz;  $\text{CDCl}_3$ ): d 4.32 (1H, br); 4.08 (5H, br); 4.02 (1H, br); 3.94 (1H br); 2.84 (4H, br); 1.8-2.5 (8H, br); 1.05-1.4 (24H, br,).

$^{31}\text{P}$  NMR (101 MHz;  $\text{CDCl}_3$ ): d 4.15 ppm.

25

Elemental analysis: Found: C:64.26%; H:7.88%  
Calculated: C:65.03%; H:7.94%

Example 24Preparation of methyl propanoate from ethylene, carbon monoxide and methanol catalysed by a compound of the present invention

5

A mechanically stirred autoclave (Hastelloy) of 2 litre capacity was evacuated of air and then charged with a solution of tri(dibenzylideneacetone)dipalladium ( $1.44 \times 10^{-5}$  moles), 1,2-bis-(di-tertbutylphosphinomethyl) ferrocene of Example 2, ( $7.61 \times 10^{-5}$  moles) and methane sulfonic acid ( $2.25 \times 10^{-3}$  moles) in 300 ml of methyl propanoate/methanol (70 wt% methyl propanoate). The autoclave was heated to 100°C and when at that temperature, ethylene ( $8 \times 10^5 \text{Nm}^{-2}$ ) was added on top of the vapour pressure of the solvents and immediately an equimolar mixture of carbon monoxide and ethylene ( $2 \times 10^5 \text{Nm}^{-2}$ ) added to the system through a pressure regulating valve set to  $10 \times 10^5 \text{Nm}^{-2}$  above the solvent vapour pressure. Suitably, the molar ratio of ethylene to carbon monoxide in the reactor is approximately 9:1. The temperature of the reactor was maintained at 100°C and as the reaction proceeded additional carbon monoxide and ethylene was added (on an equimolar basis) through the pressure regulating Tescom valve. No catalyst precipitation was observed.

Initial reaction rates measured in moles of methyl propanoate (MeP) per mole of palladium per hour and turnover measured in moles of methyl propanoate per mole of palladium were determined for the catalyst. This may be accomplished by an analysis of the amount of gas consumed per unit time (rate) and the total amount of gas consumed

during the reaction, assuming ideal gas behaviour and 100% selectivity to methyl propanoate.

The reaction was repeated (Run 2) and initial reaction rates and turnover numbers calculated as described above. The data for both runs is displayed in Table 1.

	Maximum (moles MeP/mole Pd/hr) Initial Rate	Turnover (moles MeP/mole Pd/hr) number after 3 hours
Run 1	31,810	59,941
Run 2	30,322	63,941

Table 1

10

Comparative Example 25

Preparation of methyl propanoate from ethylene, carbon monoxide and methanol catalysed by a known catalyst

15 Example 24 was repeated twice (Runs 3 and 4 respectively) except the catalyst system was as disclosed in WO 96/19434 and obtained by charging the autoclave with tri(dibenzylideneacetone)dipalladium ( $1.44 \times 10^{-5}$  moles), 1,2 bis(di-t-butylphosphinomethyl)benzene ( $7.61 \times 10^{-5}$  moles) and methane sulfonic acid ( $2.25 \times 10^{-3}$  moles) in 20 methyl propanoate/methanol (300 ml, 70 wt% methyl propanoate).

The initial reaction rates (moles MeP/mole Pd per hour) and turnover numbers (moles Pd/moles MeP) for the catalyst were calculated as described in Example 24 above. The results are presented in Table 2 below.

	Maximum Initial Rate (moles Pd/moles MeP/hr)	Turnover number after 3 hours (moles Pd/mole MeP)
Run 3	29,730	48,386
Run 4	30,335	51,997

Table 2

5 The results demonstrate (see Table 1 and Table 2), that the catalyst of the present invention palladium 1,2-bis-(di-t-butylphosphinomethyl)ferrocene and the known palladium 1,2-bis(di-t-butylphosphinomethyl)benzene exhibit comparable initial catalytic reaction rates.

10 However, the turnover number for the catalyst of the present invention is significantly higher than that for the known palladium 1,2-bis(di-t-butylphosphinomethyl)benzene catalyst, thereby indicating that the compound of the present invention increases the

15 rate of the carbonylation reaction compared to the known bidentate system.

Example 26

Preparation of methyl propanoate from ethylene, carbon

20 monoxide and methanol catalysed by a compound of the  
present invention

Example 24 was repeated (Runs 5 to 9 respectively) except the catalyst system employed was obtained by charging the

25 autoclave with 1,2-bis-(di-1-adamantylphosphinomethyl)ferrocene-bis-methane sulphonate (Example 13 or 14,  $7.61 \times 10^{-5}$  moles), tri(dibenzylideneacetone)dipalladium ( $1.44 \times 10^{-5}$  moles) and methane sulfonic acid ( $2.25 \times 10^{-3}$  moles)

in methyl propanoate/methanol (300 ml, 70 wt% methyl propanoate).

The initial reaction rates (moles MeP/mole Pd per hour) and turnover numbers (moles Pd/moles MeP) for the catalyst were calculated as described in Example 24 above. The results are presented in Table 3 below.

	Maximum Initial Rate (moles Pd/moles MeP/hr)	Turnover number after 3 hours (moles Pd/mole MeP)
Run 5	52,854	67,885
Run 6	37,034	64,996
Run 7	35,986	64,441
Run 8	40,781	62,108
Run 9	39,251	62,108


Table 3

The results demonstrate (see Table 3 and Table 2), that the catalyst of the present invention palladium 1,2-bis(di-1-adamantylphosphinomethyl)ferrocene-bis-methane sulphonate exhibits a significantly higher initial catalytic rate and a significantly higher turnover number than the known catalyst palladium 1,2-bis-(di-t-butylphosphinomethyl)benzene, thereby indicating the compound of the present invention increases the rate of the carbonylation reaction compared to the known bidentate system.

# Claims

1. A compound obtainable by combining:

- (a) a Group VIIIIB metal or a compound thereof; and,  
 5 (b) a compound of formula I or salt thereof:



wherein:

10  $A_1$  and  $A_2$ , and  $A_3$ ,  $A_4$  and  $A_5$  (when present), each independently represent lower alkylene;

K is selected from the group consisting of hydrogen, lower alkyl, aryl, Het, halo, cyano, nitro,  $-OR^{19}$ ,  $-OC(O)R^{20}$ ,  
 15  $-C(O)R^{21}$ ,  $-C(O)OR^{22}$ ,  $-N(R^{23})R^{24}$ ,  $-C(O)N(R^{25})R^{26}$ ,  $-C(S)(R^{27})R^{28}$ ,  
 $-SR^{29}$ ,  $-C(O)SR^{30}$ ,  $-CF_3$  or  $-A_3-Q^3(X^5)X^6$ ;

D is selected from the group consisting of hydrogen, lower alkyl, aryl, Het, halo, cyano, nitro,  $-OR^{19}$ ,  $-OC(O)R^{20}$ ,  
 20  $-C(O)R^{21}$ ,  $-C(O)OR^{22}$ ,  $-N(R^{23})R^{24}$ ,  $-C(O)N(R^{25})R^{26}$ ,  $-C(S)(R^{27})R^{28}$ ,  
 $-SR^{29}$ ,  $-C(O)SR^{30}$ ,  $-CF_3$  or  $-A_4-Q^4(X^7)X^8$ ;

E is selected from the group consisting of hydrogen, lower alkyl, aryl, Het, halo, cyano, nitro,  $-OR^{19}$ ,  $-OC(O)R^{20}$ ,



$-\text{C}(\text{O})\text{R}^{21}$ ,  $-\text{C}(\text{O})\text{OR}^{22}$ ,  $-\text{N}(\text{R}^{23})\text{R}^{24}$ ,  $-\text{C}(\text{O})\text{N}(\text{R}^{25})\text{R}^{26}$ ,  $-\text{C}(\text{S})(\text{R}^{27})\text{R}^{28}$ ,  
 $-\text{SR}^{29}$ ,  $-\text{C}(\text{O})\text{SR}^{30}$ ,  $-\text{CF}_3$  or  $-\text{A}_5-\text{Q}^5(\text{X}^9)\text{X}^{10}$ ;

or both D and E together with the carbon atoms of the  
 5 cyclopentadienyl ring to which they are attached form an  
 optionally substituted phenyl ring:

$\text{X}^1$  represents  $\text{CR}^1(\text{R}^2)(\text{R}^3)$ , congressyl or adamantyl,  $\text{X}^2$   
 represents  $\text{CR}^4(\text{R}^5)(\text{R}^6)$ , congressyl or adamantyl, or  $\text{X}^1$  and  
 10  $\text{X}^2$  together with  $\text{Q}^2$  to which they are attached form an  
 optionally substituted 2-phospha-adamantyl group, or  $\text{X}^1$   
 and  $\text{X}^2$  together with  $\text{Q}^2$  to which they are attached form a  
 ring system of formula 1a;

15  $\text{X}^3$  represents  $\text{CR}^7(\text{R}^8)(\text{R}^9)$ , congressyl or adamantyl,  $\text{X}^4$   
 represents  $\text{CR}^{10}(\text{R}^{11})(\text{R}^{12})$ , congressyl or adamantyl, or  $\text{X}^3$   
 and  $\text{X}^4$  together with  $\text{Q}^1$  to which they are attached form an  
 optionally substituted 2-phospha-adamantyl group, or  $\text{X}^3$   
 and  $\text{X}^4$  together with  $\text{Q}^1$  to which they are attached form a  
 20 ring system of formula 1b;

$\text{X}^5$  represents  $\text{CR}^{13}(\text{R}^{14})(\text{R}^{15})$ , congressyl or adamantyl,  $\text{X}^6$   
 represents  $\text{CR}^{16}(\text{R}^{17})(\text{R}^{18})$ , congressyl or adamantyl, or  $\text{X}^5$   
 and  $\text{X}^6$  together with  $\text{Q}^3$  to which they are attached form an  
 25 optionally substituted 2-phospha-adamantyl group, or  $\text{X}^5$   
 and  $\text{X}^6$  together with  $\text{Q}^3$  to which they are attached form a  
 ring system of formula 1c;

$\text{X}^7$  represents  $\text{CR}^{31}(\text{R}^{32})(\text{R}^{33})$ , congressyl or adamantyl,  $\text{X}^8$   
 30 represents  $\text{CR}^{34}(\text{R}^{35})(\text{R}^{36})$ , congressyl or adamantyl, or  $\text{X}^7$   
 and  $\text{X}^8$  together with  $\text{Q}^4$  to which they are attached form an  
 optionally substituted 2-phospha-adamantyl group, or  $\text{X}^7$

and  $X^8$  together with  $Q^4$  to which they are attached form a ring system of formula 1d;

$X^9$  represents  $CR^{37}(R^{38})(R^{39})$ , congressyl or adamantyl,  $X^{10}$   
5 represents  $CR^{40}(R^{41})(R^{42})$ , congressyl or adamantyl, or  $X^9$   
and  $X^{10}$  together with  $Q^5$  to which they are attached form an  
optionally substituted 2-phospha-adamantyl group, or  $X^9$   
and  $X^{10}$  together with  $Q^5$  to which they are attached form a  
ring system of formula 1e;

10

$Q^1$  and  $Q^2$ , and  $Q^3$ ,  $Q^4$  and  $Q^5$  (when present), each  
independently represent phosphorus, arsenic or antimony;

M represents a Group VIB or VIIIB metal or metal cation  
15 thereof;


$L_1$  represents an optionally substituted cyclopentadienyl,  
indenyl or aryl group;

20  $L_2$  represents one or more ligands each of which are  
independently selected from hydrogen, lower alkyl,  
alkylaryl, halo, CO,  $P(R^{43})(R^{44})R^{45}$  or  $N(R^{46})(R^{47})R^{48}$ ;

$R^1$  to  $R^{18}$  and  $R^{31}$  to  $R^{42}$ , when present, each independently  
25 represent hydrogen, lower alkyl, aryl, halo or Het;

$R^{19}$  to  $R^{30}$  and  $R^{43}$  to  $R^{48}$ , when present, each independently  
represent hydrogen, lower alkyl, aryl or Het;

30 the ring systems of formula 1a, 1b, 1c, 1d and 1e are  
represented by the formulae



$R^{49}$ ,  $R^{54}$  and  $R^{55}$ , each independently represent hydrogen, lower alkyl or aryl;  $R^{50}$  to  $R^{53}$  each independently represent hydrogen, lower alkyl, aryl or Het; and  $Y^1$ ,  $Y^2$ ,  $Y^3$ ,  $Y^4$  and  $Y^5$ , each independently represent oxygen, sulfur or  $N-R^{55}$ ;

$n = 0$  or  $1$ ;

and  $m = 0$  to  $5$ ;

provided that when  $n = 1$  then  $m$  equals  $0$ , and when  $n$  equals  $0$  then  $m$  does not equal  $0$ .

2. A compound as claimed in claim 1, wherein if both  $K$  represents  $-A_3-Q^3(X^5)X^6$  and  $E$  represents  $-A_5-Q^5(X^9)X^{10}$ , then  $D$  represents  $-A_4-Q^4(X^7)X^8$ .

3. A compound as claimed in claim 1 or 2, wherein  $R^1$  to  $R^{18}$  and  $R^{31}$  to  $R^{42}$  each independently represent hydrogen, optionally substituted  $C_1$ - $C_6$  alkyl or optionally substituted phenyl.

4. A compound as claimed in any one of claims 1 to 3, wherein  $R^1$  to  $R^{18}$  and  $R^{31}$  to  $R^{42}$  each independently represent hydrogen or non-substituted  $C_1$ - $C_6$  alkyl.

5. A compound as claimed in any one of claims 1 to 3, wherein one or more of the groups  $R^1$  to  $R^3$ ,  $R^4$  to  $R^6$ ,  $R^7$  to

R<sup>9</sup>, R<sup>10</sup> to R<sup>12</sup>, R<sup>13</sup> to R<sup>15</sup>, R<sup>16</sup> to R<sup>18</sup>, R<sup>31</sup> to R<sup>33</sup>, R<sup>34</sup> to R<sup>36</sup>, R<sup>37</sup> to R<sup>39</sup>, R<sup>40</sup> to R<sup>42</sup> together with the carbon atom to which they are attached each independently form a cyclic alkyl structure.

5

6. A compound as claimed in any one of claims 1 to 3, wherein one or more of the groups R<sup>1</sup> and R<sup>2</sup>, R<sup>4</sup> and R<sup>5</sup>, R<sup>7</sup> and R<sup>8</sup>, R<sup>10</sup> and R<sup>11</sup>, R<sup>13</sup> and R<sup>14</sup>, R<sup>16</sup> and R<sup>17</sup>, R<sup>31</sup> and R<sup>32</sup>, R<sup>34</sup> and R<sup>35</sup>, R<sup>37</sup> and R<sup>38</sup>, R<sup>40</sup> and R<sup>41</sup> together with the carbon atom to which they are attached each independently form a cyclic alkyl structure.

10

7. A compound as claimed in any one of the preceding claims, wherein each of R<sup>1</sup> to R<sup>18</sup> and R<sup>31</sup> to R<sup>42</sup> does not represent hydrogen.

15

8. A compound as claimed in any one of the preceding claims, wherein adamantyl represents unsubstituted adamantyl or adamantyl substituted with one or more unsubstituted C<sub>1</sub>-C<sub>8</sub> alkyl substituents, or a combination thereof.

20

9. A compound as claimed in any one of the preceding claims, wherein 2-phospha-adamantyl represents unsubstituted 2-phospha-adamantyl or 2-phospha-adamantyl substituted with one or more unsubstituted C<sub>1</sub>-C<sub>8</sub> alkyl substituents, or a combination thereof.

25

10. A compound as claimed in any one of the preceding claims, wherein 2-phospha-adamantyl includes one or more oxygen atoms in the 2-phospha-adamantyl skeleton.

30

11. A compound as claimed in any one of the preceding claims, wherein congressyl represents unsubstituted congressyl.

5 12. A compound as claimed in any one of the preceding claims, wherein  $R^{50}$  to  $R^{53}$  each independently represent optionally substituted  $C_1$ - $C_6$  alkyl, trifluoromethyl or phenyl optionally substituted with non-substituted  $C_1$ - $C_6$  alkyl or  $OR^{19}$  where  $R^{19}$  represents non-substituted  $C_1$ - $C_6$   
10 alkyl.

13. A compound as claimed in any one of the preceding claims, wherein  $R^{49}$  and  $R^{54}$  each independently represent hydrogen or non-substituted  $C_1$ - $C_6$  alkyl.

15

14. A compound as claimed in any one of the preceding claims, wherein each of  $Y^1$  to  $Y^5$  represents oxygen.

15. A compound as claimed in any one of the preceding  
20 claims, wherein  $X^1$  is identical to  $X^3$ , and  $X^5$ ,  $X^7$  and  $X^9$  when present.

16. A compound as claimed in any one of the preceding claims, wherein  $X^2$  is identical to  $X^4$ , and  $X^6$ ,  $X^8$  and  $X^{10}$   
25 when present.

17. A compound as claimed in any one of claims 1 to 14, wherein  $X^1$  represents  $CR^1(R^2)(R^3)$ ,  $X^2$  represents  $CR^4(R^5)(R^6)$ ,  $X^3$  represents  $CR^7(R^8)(R^9)$  and  $X^4$  represents  
30  $CR^{10}(R^{11})(R^{12})$ .

18. A compound as claimed in any one of claims 1 to 14, wherein  $X^1$  represents  $CR^1(R^2)(R^3)$ ,  $X^2$  represents adamantyl,  $X^3$  represents  $CR^7(R^8)(R^9)$  and  $X^4$  represents adamantyl.

5 19. A compound as claimed in any one of claims 1 to 14, wherein  $X^1$  represents  $CR^1(R^2)(R^3)$ ,  $X^2$  represents congressyl,  $X^3$  represents  $CR^7(R^8)(R^9)$  and  $X^4$  represents congressyl.

10 20. A compound as claimed in any one of claims 1 to 14, wherein  $X^1$  to  $X^4$  each independently represent adamantyl.

21. A compound as claimed in any one of claims 1 to 14, wherein  $X^1$  to  $X^4$  each independently represent congressyl.

15

22. A compound as claimed in any one of claims 1 to 14, wherein  $X^1$  and  $X^2$  together with  $Q^2$  to which they are attached form a ring system of formula Ia, and  $X^3$  and  $X^4$  together with  $Q^1$  to which they are attached form a ring  
20 system of formula Ib.

23. A compound as claimed in any one of claims 1 to 14, wherein  $X^1$  and  $X^2$  together with  $Q^2$  to which they are attached form a 2-phospha-adamantyl group, and  $X^3$  and  $X^4$   
25 together with  $Q^1$  to which they are attached form a 2-phospha-adamantyl group.

24. A compound as claimed in any one of the preceding claims, wherein K represents hydrogen.

30

25. A compound as claimed in any one of claims 1 to 23, wherein K represents  $-A_3-Q^3(X^5)X^6$ .

26. A compound as claimed in claim 25, wherein  $-A_3-Q^3(X^5)X^6$  is identical to  $-A_2-Q^1(X^3)X^4$ .

27. A compound as claimed in any one of the preceding  
5 claims, wherein D and E together with the carbon atoms of the cyclopentadienyl ring to which they are attached form an unsubstituted phenyl ring.

28. A compound as claimed in any one of the preceding  
10 claims, wherein D and E both represent hydrogen.

29. A compound as claimed in any one of claims 1 to 26, wherein D represents  $-A_4-Q^4(X^7)X^8$ .

15 30. A compound as claimed in claim 29, wherein  $-A_4-Q^4(X^7)X^8$  is identical to  $-A_2-Q^1(X^3)X^4$ .

31. A compound as claimed in any one claims 29 or 30, wherein E represents hydrogen.

20

32. A compound as claimed in any one claims 1 to 26, 29 or 30, wherein E represents  $-A_5-Q^5(X^9)X^{10}$ .

33. A compound as claimed in claim 32, wherein  $-A_5-Q^5(X^9)X^{10}$  is identical to  $-A_2-Q^1(X^3)X^4$ .

25

34. A compound as claimed in any one of the preceding claims, wherein  $A_1$  and  $A_2$ , and  $A_3$ ,  $A_4$  and  $A_5$  when present, each independently represent  $-CH_2-$  or  $-C_2H_4-$ .

30

35. A compound as claimed in any one of the preceding claims, wherein each  $A_1$  and  $A_2$ , and  $A_3$ ,  $A_4$  and  $A_5$  when present are identical and preferably represent  $-CH_2-$ .

36. A compound as claimed in any one of the preceding claims, wherein each  $Q^1$  and  $Q^2$ , and  $Q^3$ ,  $Q^4$  and  $Q^5$  when present are identical and preferably represent  
5 phosphorous.

37. A compound as claimed in any one of the preceding claims, wherein  $n=1$ ,  $m=0$  and  $L_1$  is selected from cyclopentadienyl, phenyl, indenyl or naphthyl, preferably  
10 unsubstituted cyclopentadienyl.


38. A compound as claimed in any one of the preceding claims, wherein  $M$  represents iron or a metal cation thereof.  
15

39. A compound as claimed in any one of the preceding claims obtainable by combining: (a) palladium or a compound thereof; and (b) a compound of formula I as defined in any one of the preceding claims.  
20

40. A process for preparing a compound as defined in any one of claims 1 to 39 comprising combining (a) a Group VIIIB metal or compound thereof; and, (b) a compound of formula I as defined in any one of claims 1 to 38.  
25

41. A compound of formula I







wherein  $A_1$ ,  $A_2$ ,  $K$ ,  $D$ ,  $E$ ,  $M$ ,  $L_2$ ,  $L_1$ ,  $Q^1$ ,  $Q^2$ ,  $X^1$ ,  $X^2$ ,  $X^3$ ,  $X^4$ ,  $n$  and  $m$  are as defined in any one of claims 1 to 38.

5


42. A process for preparing a compound of formula I as defined in claim 41, comprising reacting a compound of formula II wherein  $A_1$ ,  $A_2$ ,  $K$ ,  $D$ ,  $E$ ,  $M$ ,  $L_1$ ,  $L_2$ ,  $n$  and  $m$  are as defined for a compound of formula I, and  $LG_1$  and  $LG_2$  represent suitable leaving groups, with a compound of formula IIIa and IIIb

10





(IIIa)




(IIIb)

5 wherein  $\text{X}^1$ ,  $\text{X}^2$ ,  $\text{Q}^2$ ,  $\text{X}^3$ ,  $\text{X}^4$  and  $\text{Q}^1$  are as defined in anyone of claims 1 to 38.


43. A compound of formula II as defined in claim 42.

10 44. A process for preparing a compound of formula I wherein K, D, E, M,  $\text{A}_2$ ,  $\text{A}_1$ ,  $\text{L}_2$ ,  $\text{L}_1$ ,  $\text{Q}^1$ ,  $\text{Q}^2$ , m and n are as defined in any one of claims 1 to 38 and  $\text{X}^1$  and  $\text{X}^2$  together with  $\text{Q}^2$  to which they are attached form a ring system of formula Ia as defined in anyone of claims 1 to  
 15 38 and  $\text{X}^3$  and  $\text{X}^4$  together with  $\text{Q}^1$  to which they are attached form a ring system of formula Ib as defined in any one of claims 1 to 38, comprising reacting a compound of formula XV



20

wherein K, D, E, M,  $\text{A}_2$ ,  $\text{A}_1$ ,  $\text{L}_2$ ,  $\text{L}_1$ ,  $\text{Q}^1$ ,  $\text{Q}^2$ , m and n are as defined in any one of claims 1 to 38, with a compound of formula XVIa and XVIb



wherein Y<sup>1</sup>, Y<sup>2</sup>, R<sup>49</sup> to R<sup>55</sup> are as defined for a compound of formula I.

5

45. A compound of formula XV as defined in claim 44.

46. A process for the carbonylation of an ethylenically unsaturated compound comprising contacting an ethylenically unsaturated compound with carbon monoxide and a co-reactant in the presence of a compound as defined in any one of claims 1 to 39.

47. A process as defined in claim 46 wherein the co-reactant includes a hydroxyl group containing compound.

48. A process as claimed in claim 46 or 47 wherein the ethylenically unsaturated compound comprises ethylene, 1,3-butadiene, oct-1-ene or vinyl acetate, preferably ethylene.

20

49. A process as claimed in any one of claims 46 to 48, further including the step of including a source of anions.

25

50. A composition comprising a compound as defined in any one of claims 1 to 39 attached to a support.

51. Use of a compound as defined in anyone of claims 1 to 39 or a composition as defined in claim 50 as a catalyst.